

Review for Test #3  
Neutron Stars and Black Holes

---

Neutron stars – mass of sun, radius ~ 10km, density like atomic nucleus, huge gravity at surface.

Discovery of pulsars – pulsating radio sources

Interpretation of pulsars as rotating magnetized neutron stars

Role of magnetic field to cause radiation, misalignment of rotation axis, magnetic axis

Production of pulses – probably related to strong electric, magnetic fields at magnetic poles

About 600 pulsars known, perhaps a billion neutron stars in the Galaxy.

Pressure support from quantum pressure of neutrons plus nuclear repulsion. Maximum mass of neutron star about 2 solar masses.

Neutron stars as binary X-rays sources.

X-ray pulsars – accreted gas channeled to magnetic poles, “pulsar” by lighthouse effect if magnetic axis is tilted with respect to the spin axis.

X-ray transients – 4 or 5 known in our Galaxy. Outburst every few years for a month. Probably a disk instability like a dwarf nova, but with the white dwarf replaced by a neutron star.

X-ray Bursters – about 30 known in our Galaxy. Burst every few hours for minutes. Probably the neutron star analog of a classical nova. Matter accretes on surface of neutron star. Hydrogen is supported by thermal pressure, burns to helium. Helium is supported by quantum pressure and is unregulated and explodes, producing the X-rays.

Magnetars – neutron stars with magnetic fields 100 to 1000 times stronger than the Crab nebula pulsar.

Soft gamma-ray repeaters – objects that emit intense bursts of low energy gamma rays and X-rays for a few minutes every few years. Periodic “pulses” after the initial flash. Observed spin-down rates imply they are magnetars. One soft gamma-ray repeater actually caused aurorae and interfered with terrestrial radio communications August, 1998, another flared on the far side of our Galaxy, and was detected on December 27, 2004.

Black Hole History – Mitchell, Laplace, escape velocity.

Conceptual problems with Newton’s Theory of Gravity

Einstein says there is no “force” of gravity. Matter curves space and curved space tells matter how to move.

Dimension – determined by the number of mutually perpendicular directions in a given space

Space versus Hyperspace

Black Hole Evaporation—For a black hole of ordinary stellar mass or larger the amount of mass loss is negligible in the age of the Universe and may be ignored. A black hole of less than asteroid mass could totally evaporate within the age of the Universe.

The three fundamental properties of a Black Hole are those that can be measured from a distance - mass, charge, and spin. Other properties such as size and shape are specified once these basic properties are set.

Information Loss in Black Holes – Quantum theory insists information is preserved, black holes seem to destroy it.

Time-like space—interior to event horizon space drags in one direction, just as time drags you older.

Schwarzschild black hole—mass but no spin, no electrical charge. Time-like space leads to the singularity, so it cannot be avoided.

Rotating or Kerr black hole—the idealized mathematical solution of Einstein's equations developed by Kerr in which one assumes that all the mass is in the rotating singularity and that there is vacuum everywhere else.

Singularity in a rotating black hole—shaped like a ring, surrounded by “normal ” space so that it can be avoided in principle.

Time-like space in rotating black hole—the “in-going” time-like space is bounded on both sides by an event horizon so that it does not extend down to the singularity. Inside the inner event horizon is “normal” space surrounding the singularity. At the same place, but in the future, there is a region of “out-going” time-like space again bounded by two event horizons leading out to a normal Universe of flat space. In the future of that Universe is another in-going time-like space.

Inner “normal” space. Inside the rotating black hole the “normal” space will be one of huge gravity and tidal forces, but they are not infinite, and one could survive in principle never emerging from the black hole, but also never hitting the singularity.

Through the singularity—passing through the ring of the singularity leads to another volume of “normal” space within the black hole surrounding the singularity, but it is not the same one that surrounds the singularity that is first encountered when entering the black hole.

Blue shift - in a real Universe matter and energy falling into a black hole will gain energy (blue shift) and that energy will probably alter the “vacuum” Kerr solution, so no extra Universes are accessible.

Clues for black holes – look for binary system where X-rays are produced in accretion disk before matter disappears down the black hole and Kepler’s law helps to determine mass greater than maximum mass of neutron star.

Cygnus X-1 – First candidate black hole in a binary star system. Object of  $10M_{\odot}$  emits X-rays and orbits unevolved star of  $30M_{\odot}$ . Small probability that  $10M_{\odot}$  object is itself a  $9M_{\odot}$  star transferring mass to a  $1M_{\odot}$  neutron star. The  $9M_{\odot}$  star could be lost in glare of  $30M_{\odot}$  star.

Black hole candidates with low mass companion stars – for these systems the “unseen” X-ray emitting star is more massive than the unevolved companion. No third ordinary star could remain unseen.

Black hole X-ray novae-all recently discovered black hole candidates sit undiscovered for decades then flare for a few months. Thought to be flushing instability in accretion disk, occurs in systems with low mass unevolved companions.