

February 22, 2010

Second exam, Friday. Review Sheet Today. Review Session Thursday, Room WEL 2.312. (NOT RLM 15.216B).


Reading, Sections 6.4, 6.5, 6.6, 6.7

Sections 1.2, 2.1, 2.4, 2.5, Sections 3.3, 3.4, 3.5, 3.10, Sections 4.1, 4.2, 4.3, 4.4, Sections 5.2, 5.4 for background

Astronomy in the News? Space shuttle returned in night landing.


Some pushback in Congress over proposal to cut back the Constellation Moon program

Pic of the Day - A group of interacting galaxies



Goal - understand how accretion disks work, what sort of radiation they emit.

Basic Disk Dynamics - Figure 4.1



Basic Disk Dynamics

Orbits closer to the center are faster.

This creates rubbing and friction and heat, everywhere in the disk.

Friction tries to slow the orbiting matter, but it falls *inward* and ends up moving *faster*.


(Just as removing heat from a normal star causes it to get hotter)

Slow settling inward by friction -- *accretion*

Friction also causes *heat*.

Hotter on inside, cooler on outside

Optical → UV → X-rays



One Minute Exam:

In an accretion disk, friction causes moving matter to



Slow down



Speed up



Move outward



Pass from one Roche lobe to another

Goal - understand how white dwarfs in binary star systems can, and cannot grow to the Chandrasekar mass and explode.


Cataclysmic Variables

Second stage of mass transfer

General Category “Novae”

“New” stars flare up, see where none had been seen before.

All CVs share same general features: *transferring star*, *transfer stream*, *hot spot*, *accretion disk*, and *white dwarf*.



Classical Novae: Problem with losing mass from white dwarf during surface explosions.

Recurrent Novae like USco: do seem to have large mass white dwarfs, encouraging, but maybe not enough.

§ 5.4 Final Evolution of Cataclysmic Variables

Some CVs have managed to reach large masses

$M_{\text{wd}} \sim M_{\text{ch}}$ Chandrasekhar mass, 1.4 solar masses, like U Sco

If get close enough to M_{ch} , attain high density,
ignite carbon in center

Quantum Deregulated \rightarrow violent explosion

Type Ia Supernova?!

What CVs have white dwarfs that reach M_{ch} ?

Not classical novae

explosion of surface H shell also rips off a bit of the
white dwarf mass - we see excess carbon & oxygen in
ejected matter

white dwarf shrinks in mass rather than grows.

Likely outcome in this case - 2nd star finally burns out H, tries to form
red giant, mass transfer \Rightarrow ***Two WDs!***

Sky Watch

Classical Novae:

CP Pup, toward constellation Puppis in 1942

Pup 91, another toward Puppis in 1991 (not same place in our Galaxy, just accidentally off in the same approximate direction)

QU Vul, toward constellation Vulpecula, white dwarf composed of Oxygen, Neon, and Magnesium rather than Carbon and Oxygen.

GK Per toward constellation Perseus - has had both a classical nova eruption in 1901 and dwarf nova eruptions.

Sky Watch

Recurrent Novae:





U Sco in the constellation Scorpius is a Recurrent Nova,
It may be a candidate to explode as a supernova!

Might see Scorpius. Also has neutron stars and black holes.

T Pyx in constellation Pyxis.

One Minute Exam

We expect classical nova systems to end up making two white dwarfs orbiting one another because:

-  The first white dwarf loses mass and hence cannot grow and explode
-  The first white dwarf will accrete mass until it reaches the Chandrasekhar limit
-  The main sequence star transferring mass must eventually make a white dwarf
-  The second white dwarf has the Chandrasekhar mass

Goal - understand what happens to two white dwarfs in a binary system.


We do observe 2 white dwarfs in orbit in some cases - is that the end?

No: *gravitational radiation* (§ 3.10)

ripples in curved space-time

like paddle on surface of pond

remove energy from orbit - acts as drag



If you try to slow down an orbiting object what happens?


Falls inward, speeds up,

Get more gravitational radiation, more inspiral

Given enough time (billions of years) 2 white dwarfs must spiral together!

What happens when two white dwarfs spiral together?

New physical fact:
Larger mass WD has smaller radius



Which WD has the smaller Roche lobe?

Which fills its Roche Lobe first?

When the first WD fills its Roche lobe,
what happens to its radius?

When the first WD fills its Roche lobe,
what happens to its Roche lobe?

What happens to the white dwarf?