

April 5, 2010

Reading: Chapter 10, Sections 10.1-10.4, 10.9

Wheeler on travel Wednesday and Friday, No Wheeler office hours today (or Wed, Fri).

Exam 4, April 16.

Astronomy in the News?

Pic of the Day - Saturn “shepherd” moon,
Prometheus



Chapter 10 - Finding Black Holes for Real

We know that massive stars evolve to form iron cores that absorb energy and collapse. A compact object must be left behind.

Some explode and leave rotating, magnetic pulsars

Some explode and leave highly magnetic magnetars

Some explode but leave black holes or completely collapse to leave black holes

We don't know which massive stars do which! Tendency to think that more massive stars are more prone to making black holes, but the rotation of the star, the presence of a binary companion, and other factors may influence the outcome.

We do know that black holes exist, so some stars make them.

Goal:

To understand how we search for real black holes and why binary systems with mass transfer and accretion disks are so important.

Black Holes for Real

There may be 1 - 100 million black holes in the Galaxy made by collapsing stars over the history of the Galaxy.

That means that the nearest black hole may be only a few tens of light years away. How do we find them?

Black holes made from stars are really black! (Negligible Hawking radiation).

Those alone in space are not impossible to find, but very tough. None yet identified.


Look for binary systems, where mass accretion occurs.

Will not see the black hole, cannot yet “see” a black spot.

Can detect the *halo of X-rays* from orbiting matter, the accretion disk, near the event horizon that will reveal the presence and nature of the black hole. ***Look in accreting binary systems!***



Goal is to get close-up study of strongly warped space



Perez and Wagoner, Stanford: computer simulation of radiation from inner black hole accretion disk

Black holes are so weird and so important that the standards of evidence have to be high!

Current evidence is still primarily circumstantial, but very strong:

Stellar mass black holes (several to ~ 10 solar masses), binary systems in our Galaxy or nearby galaxies

Intermediate mass black holes ($\sim 1000 - 10,000$ solar masses)??, in binary systems or stellar clusters in our Galaxy or nearby galaxies

Supermassive black holes (million to a billion solar masses) in the middle of our Galaxy and in the middle of many, many others.

Circumstantial arguments for presence of black hole in a binary system:

Only neutron stars and black holes have the high gravity necessary for intense X-rays.

Use Kepler's laws to measure the total mass of the system, astronomy to determine the mass of the mass-losing star, subtract to get mass of "unseen" companion emitting X-rays.

Maximum mass of neutron star is ~ 2 solar masses

Intense X-ray source with mass exceeding 2 solar masses is, by a process of elimination, a candidate black hole.

There are about 20 binary star black hole candidates in our Galaxy and in the Large Magellanic Cloud (near enough to detect the X-rays) that have masses measured to be greater than 3 solar masses, and hence too massive to be a neutron star.

There are another 25 binary star black hole candidates with similar X-ray properties, but no measured mass.

Cygnus X-1

First X-ray source discovered in the direction of the constellation Cygnus.

Discovered in 1970's by Uhuru Satellite (Swahili for Freedom).


First and still most famous stellar-mass binary black hole candidate.

Can't see this system with the naked eye, but can find constellation Cygnus - look for it for sky watch!


Cygnus X-1

Optically dark
X-ray emitting companion
 $\geq 10M_{\odot} \gg$ NS max mass
 \Rightarrow BH


Could nature be
tricking us? All
we really know
is that there is a
 $10M_{\odot}$ “thing”
emitting X-rays



$M \sim 10M_{\odot}$
Not NS



Blue supergiant, mass
losing star



One possibility:
 $9M_{\odot}$ normal star
“lost in glare” of $30M_{\odot}$
like flashlight next to
searchlight. Unlikely but
can't be entirely ruled out.


Expect only two or three systems like Cygnus X-1 in our Galaxy.

Bright, massive, short-lived companion

Maybe only one, and we found it!

Surprisingly, most binary black hole candidates have small mass main sequence companions, typically $\sim 1/2$ solar mass.

Observe ~ 45 such systems and guess there may be ~ 1000 in the Galaxy



For systems with small mass companions cannot hide a 3rd star in the system

⇒ best black hole candidates.

Evidence still circumstantial but virtual proof of black hole

Candidates in the directions of Sagittarius, Ursa Majoris, Perseus, Scorpius, Ophiuchus, Vulpecula, Monoceros, Lupus, Cygnus (2) (Find and observe the constellations for extra credit)

AO620-00 = Nova Mon 1975 = V616 Monocerotis - one of the first and best studied with a small mass companion, black hole about 5 solar masses.

V404 Cygni - somewhat evolved companion, but one of the best cases for a black hole with “dark” mass of about 12 solar masses.

Two candidates in the Large Magellanic Cloud:
LMC X-1, LMC X-3

Total number of such systems known, about 45.

Not sure how these binary systems form.

Would have expected massive stars that can make black holes in core collapse to have massive companions, like Cygnus X-1.

Need to have black hole very close to small mass companion, current separation smaller than size of the star that made the black hole.

Possibilities:

Black hole progenitor swallows small mass companion while a red giant?

Companion forms from left-overs of collapse?

Proving Black Holes

Astronomers search for ways to directly determine that the dark X-ray producing object is a black hole, not a neutron star.


How would you identify a black hole of 1 solar mass?


Evidence that in some circumstances black holes, but not neutron stars, can produce very hot, rarified inner accretion regions, making gamma-rays, but few X-rays.

This is evidence that the object has **no surface**.

One Minute Exam

The best candidate for a binary star system with black hole is:

 One with a 30 solar mass ordinary star

 One with a 1/2 solar mass ordinary star

 One with two black holes in orbit

 Cygnus X-1