


Cosmic Evolution, Part II
Heavy Elements to Molecules

First a review of terminology:



Neutral atom:
ion:

e.g. C^{+2}


Electrons = # protons




Carbon nucleus + 4 (6-2) electrons

Forces

H atom



H₂
Molecule



Attractive

Repulsive




Molecule: Repulsive ~ Attractive

More delicate than atoms,
can be much more complex

“Bond” is sharing of electrons

Is molecule stable?

Yes, if EM potential energy less than separate atoms





Activation energy lower $\rightarrow T \sim 100 - 1000 \text{ K}$

(Room Temperature)

Questions

- Why is room temperature around 300 K?
- How commonly is this temperature found in the Universe?

Conventions:



Maximum # of Bonds:

H 1

O 2

N 3

C 4

Carbon very versatile

→ Complex chemistry


Interstellar Molecules

Exist as gas (individual molecules)

A few known in 1930's

Many more since 1968 - Radio astronomy


Rotation



Radio Telescope



Vibration



Infrared

Infrared Telescope



Appendix 2

Interstellar Molecules

Species	Name	Species	Name
H ₂	molecular hydrogen	CO ₂	carbon dioxide
C ₂	diatomic carbon	OCS	carbonyl sulfide
CH	methylidyne	SO ₂	sulfur dioxide
CH ⁺	methylidyne ion	SiC ₂	silicon dicarbide*
CN	cyanogen	SiCN	
CO	carbon monoxide	AlCN	
CO ⁺	carbon monoxide ion	C ₂ S	
CS	carbon monosulfide	C ₂ O	dicarbon monoxide †
OH	hydroxyl	C ₃	triatomic carbon*
HCl	hydrogen chloride	MgCN	magnesium cyanide*
NH		MgNC	magnesium isocyanide*
NO	nitric oxide	NaCN	sodium cyanide*
NS	nitrogen sulfide		
SiC	silicon carbide*	C ₂ H ₂	acetylene
SiO	silicon monoxide	C ₃ H	propynylidyne (l and c)
SiS	silicon sulfide	H ₂ CO	formaldehyde
SiN	silicon nitride	H ₂ CN	
SO	sulfur monoxide	HC ₂ N	
PN		NH ₃	ammonia
CP	*	HNCO	isocyanic acid
SO ⁺	sulfoxide ion	HOCO ⁺	
NaCl	sodium chloride*	HCNH ⁺	
AlCl	aluminum chloride*	HNCS	isothiocyanic acid
KCl	potassium chloride*	C ₃ N	cynoethynyl
AlF	aluminum fluoride*†	C ₃ O	tricarbon monoxide
FeO	iron monoxide	C ₃ S	
HF		H ₂ CS	thioformaldehyde
SH		H ₃ O ⁺	hydronium ion
		SiC ₃	
H ₃ ⁺	protonated hydrogen	C ₄ H	butadiynyl
C ₂ H	ethynyl	C ₃ H ₂	cyclopropenylidene
CH ₂	methylene †	H ₂ CCC	propadienylidene
HCN	hydrogen cyanide	HCOOH	formic acid
HNC	hydrogen isocyanide	CH ₂ CO	ketene
HCO	formyl	HC ₃ N	cianoacetylene
HCO ⁺	formyl ion	HNC ₃	
HCS ⁺	thioformyl ion	CH ₂ CN	cyanomethyl
HOC ⁺	isoformyl ion †	NH ₂ CN	cyanamide
N ₂ H ⁺	protonated nitrogen	CH ₂ NH	methanimine
HNO	nitroxyl	HC ₂ NC	
H ₂ O	water	CH ₄	methane
H ₂ S	hydrogen sulfide		
H ₂ N	hydrogen nitride		
N ₂ O	nitrous oxide		

Molecular Ions

— Important Probe of conditions

173

— Discovered in Infrared
 — Discovered in UV
 — Relevant to the Origin of Life


Species	Name	Species	Name
H ₂ COH ⁺	protonated formaldehyde	HC ₅ N	cyanodiacetylene
SiH ₄	silane*	C ₇ H	
C ₄ Si	*	HCOOCH ₃	methyl formate
C ₅	pentatomic carbon*	CH ₃ C ₃ N	methylcyanoacetylene
		CH ₃ COOH	acetic acid
C ₅ H	pentynylidyne	H ₂ C ₆	
C ₅ N		CH ₂ OHCHO	glycolaldehyde
C ₂ H ₄	ethylene*		
H ₂ CCCC	butatrienylidene	CH ₃ C ₄ H	methylidiacetylene
CH ₃ OH	methanol	CH ₃ CH ₃ O	dimethyl ether
CH ₃ CN	methyl cyanide	CH ₃ CH ₂ CN	ethyl cyanide
CH ₃ NC	methyl isocyanide	CH ₃ CH ₂ OH	ethanol
CH ₃ SH	methyl mercaptan	HC ₇ N	cyanohexatriyne
NH ₂ CHO	formamide	C ₈ H	
HC ₃ HO	propynal		
HC ₃ NH ⁺		CH ₃ C ₄ CN	†
		CH ₃ CH ₃ CO	acetone
C ₆ H		NH ₂ CH ₂ COOH	glycine†
CH ₂ CHCN	vinyl cyanide	CH ₂ OHCH ₂ OH	ethylene glycol
CH ₃ C ₂ H	methylacetylene		
CH ₃ CHO	acetaldehyde	HC ₉ N	cyano-octa-tetra-yne
CH ₃ NH ₂	methylamine		
C ₂ H ₄ O	ethylene oxide	HC ₁₁ N	cyano-deca-penta-yne
CH ₂ CHOH	vinyl alcohol		

* Detected in circumstellar envelopes only
 † tentative


Look at Appendix 2
 This is an old version

Important Examples:


Water



Ammonia



Formaldehyde



Others of Note: CO Most common after H₂


HCN, HC₃N, ... HC₁₁N → Carbon chains

CH₄ (Methane)

PAHs (Polycyclic aromatic hydrocarbons)

3 Lessons

1. Complexity (Up to 13 - atoms) is extraterrestrial
May be more complex (Hard to detect)
Glycine claimed in 1994, but, so far, not confirmed
Polycyclic Aromatic Hydrocarbons (PAHs)
(Infrared evidence)
2. Dominance of Carbon
Carbon Chemistry not peculiar to Earth
3. Formation & Destruction Analogous to early Earth



Protection by dust grains: scatter and absorb ultraviolet

Dust particles

Studies of how they scatter and absorb light
(Ultraviolet → Visible → Infrared)

⇒ Two types, range of sizes up to 10^{-6} m

Carbon

PAHs → Graphite

~ Soot

Silicates

Si + O + Mg, Fe, ...

Both Produced by old stars

Formation of Interstellar Molecules

1. H_2



Must lose the potential energy difference before it falls apart ($\sim 10^{-14}$ s)

Collisions: OK in lab, too slow in space

Emit photon: very slow for H_2 (10^7 s)

$H + H + \text{catalyst} = H_2 + \text{catalyst}$

↑
surface of dust grain



Formation of Interstellar Molecules

2. More complex molecules

Problem is activation energy barrier

$T \sim 10 \text{ K} \ll \text{Barrier}$

Use reactions **without** activation energies

e.g. Molecular ions, like HCO^+

Cosmic Ray





Energy + simple mol.

→ Reactive mol.




More complex

Ion - Molecule Reactions



Molecule
or atom


Electromagnetic
Potential
Energy



No Barrier!

Separation of
Ion and Molecule

Molecules on Dust Grains



Stick on grains
"ice"


Infrared observations show this: as molecules

Vibrate, absorb infrared

e.g. H₂O absorbs at 3×10^{-6} m

CH₄ absorbs at 8×10^{-6} m

Ices on Dust Grains



Implications

1. Similar (Carbon-Dominated) Chemistry
2. Direct Role in Origin of Life?
3. Formation + Destruction analogous to Early Earth

Roles of Dust

1. Protection from UV
2. H₂ Formation
3. Freeze-out → Mantles of Ice
H₂O, NH₃, CH₄, CO₂, HCOOH, ...
↑
Methane

Star Formation

First factor in Drake Equation: The rate of star formation

Estimate of Average Star Formation Rate (R_*)

$$R_* = \frac{\text{\# of stars in galaxy}}{\text{lifetime of galaxy}} = \frac{N_*}{t_{\text{gal}}}$$

N_* : Count them? **No**

Use Gravity (Newton's Laws)

Sun orbiting center of galaxy at 270 km s^{-1} (167 miles per second)

Kinetic energy = $\frac{1}{2}$ gravitational potential energy

$$\frac{1}{2} M_{\odot} v^2 = \frac{1}{2} \frac{G M_g M_{\odot}}{R_g} \quad \leftarrow \text{Distance of Sun from center of galaxy}$$

$$\frac{R_g v^2}{G} = M_g$$

Estimate of Average Star Formation Rate (R_*)

$$(R_g = 28,000 \text{ ly}) \rightarrow M_g = 1.4 \times 10^{11} M_\odot$$

$$\text{Add mass outside Sun's orbit} \rightarrow M_g \simeq 4.6 \times 10^{11} M_\odot$$

Most is dark matter; Models indicate $8 \times 10^{10} M_\odot$ in stars

$$N_* \simeq \frac{M_g}{\text{Avg. mass of star}} = \frac{8 \times 10^{10}}{0.5} = 16 \times 10^{10}$$

$$t_{\text{gal}} \simeq 10^{10} \text{ yr} \quad (\text{studies of old stars})$$

$$R_* \simeq \frac{16 \times 10^{10} \text{ stars}}{10^{10}} = 16 \text{ stars per year}$$

Current rate: 4 stars per year

Making an Estimate

16 stars per year is an average over history of Milky Way. Current rate is about 4 stars per year. Stars formed more rapidly early in history of Milky Way. Stars at least as old as the Sun are better candidates for intelligent life.

Any number between 5 and 20 may be correct for our purposes, but understand the way we estimated it and the uncertainties.

Star Formation

Current Star Formation

Molecular Clouds

- Composition
 - H₂ (93%), He (6%)
 - Dust and other molecules (~1% by mass)
 - CO next most common after H₂, He
- Temperature about 10 K
- Density (particles per cubic cm)
 - ~100 cm⁻³ to 10⁶ cm⁻³
 - Air has about 10¹⁹ cm⁻³
 - Water about 3 x 10²² cm⁻³
- Size 1-300 ly
- Mass 1 to 10⁶ M_{sun}

A Small Molecular Cloud



ESO PR Photo 20a/99 (30 April 1999)

The "Black Cloud" B68
(VLT ANTU + FORS1)

© European Southern Observatory




Current Star Formation

- Occurs in gas with heavy elements
 - Molecules and dust keep gas cool
 - Radiate energy released by collapse
 - Stars of lower mass can form
 - Mass needed for collapse increases with T
- Star formation is ongoing in our Galaxy
 - Massive stars are short-lived
 - Star formation observed in infrared

Visible to Infrared Views



Artist's Conception



Features:
Dusty envelope
Rotation
Disk
Bipolar outflow

R. Hurt, SSC

The Protostar

- Evolution of the collapsing gas cloud
 - At first, collapsing gas stays cool
 - Dust, gas emit photons, remove energy
 - At $n \sim 10^{11} \text{ cm}^{-3}$, photons trapped
 - Gas heats up, dust destroyed, pressure rises
 - Core stops collapsing
 - The outer parts still falling in, adding mass
 - Core shrinks slowly, heats up
 - Fusion begins at $T \sim 10^7 \text{ K}$
 - Protostar becomes a main-sequence star

Summary

- Cosmic evolution builds complexity to molecules and dust
- Energy + simple things leads to complexity
- Stars form in clouds of molecules and dust
- We have estimates for the first factor in the Drake Equation, R_*
- Understand the arguments used to get this estimate