



Multi-Dimensional Simulations of Pair-Instability Supernovae

KE-JUNG CHEN^a, ALEXANDER HEGER^a & ANN ALMGREN^b


^a School of Physics and Astronomy, University of Minnesota at Twin Cities

^b Computational Research Division, Lawrence Berkeley National Lab



Introduction

Our current understanding of the formation of the first stars in the universe implies that these stars were very massive, having a typical mass scale of hundred times the mass of the sun and explode as Pair Instability Supernova (PSN). Whereas multi-dimensional simulations of most types of supernovae have been done extensively, few such simulations exist for PSN. Our goal is to understand the energetics, hydrodynamic instabilities, and nucleosynthesis of these supernovae.



(a) Collapse (b) Burn (c) Explode

The evolution of PSN

Numerical Approach


We start our simulations using one-dimensional models obtained from the KEPLERcode, spherically symmetric Lagrangian code that followed the evolution of a star up to ten seconds before maximum compression (oxygen depleted phase). Then we map the resulting one-dimensional profiles into 2D and 3D to serve as the initial conditions for CASTRO.

CASTRO

CASTRO is a new, massively parallel, multidimensional Eulerian AMR radiation-hydrodynamics code for astrophysical applications. Time integration of the hydrodynamics equations is based on a higher-order, unsplit Godunov scheme. Block-structured adaptive mesh refinement (AMR) and sub-cycling in time enable the use of high spatial resolution where it is most needed.

Mapping of Initial Model in Multi-D


Since our initial model will be in a state close to hydrostatic equilibrium, we need to be careful how we map the 1D spherically symmetric data given on a non-uniform Lagrangian grid to a multidimensional Eulerian grid. Here we present a method that numerically conserves quantities such as mass and energy that are analytically conserved in the evolution equations. Whereas this does not guarantee that the initial data will be in perfect numerical hydrostatic balance, it is at least a physically motivated constraint and is sufficient for our simulations.




Schematic for constructing a conservative density profile.

Result I


The results presented here are from 2D runs of PSN models in cylindrical symmetry where we simulated only one hemisphere.



The ²⁸Si mass fraction of 150 M_⊙ rotational PSN after evolving ~1300 secs.




The ²⁴Mg mass fraction of 150 M_⊙ PSN at ~60 secs after bounce.




The entropy of 150, 200, 250 M_⊙ PSN at 80 secs after bounce.

Result II



Figure(a) is the map of specific nuclear energy generation rate at ~60 secs after bounce. We find Rayleigh-Taylor(RT) instabilities develop at the edge of the oxygen-burning shell. Later these instabilities will grow further and affect such properties as the observable light curve of the supernova. Figure(b) is a close-up of the RT instability.



The ¹⁶O mass fraction map for rotational models at ~80 secs after bounce; they correspond to 0%, 30%, 100% of Keplerian rotational rate at radius, 2.5 x 10⁹ cm, respectively.

Summary

Discussion and Conclusion

We have presented preliminary results from our first multidimensional numerical study of the evolution of pair instability supernova using the new Eulerian AMR radiation-hydrodynamics code CASTRO. We simulated the formation of Rayleigh-Taylor instabilities in the explosion of pair instability supernovae. We have introduced a new mapping method that can be used to define the initial conditions for multidimensional simulations from one-dimensional initial data in such a way that conservation of physical quantities, monotonicity, and continuity are guaranteed at any resolution.

Acknowledgments

We would like to thank members of the Center for Computational Sciences and Engineering (CCSE) at LBNL for their support with using CASTRO. We would also like to thank Candace Joggerst and Haitao Ma for assistance with CASTRO and Laurens Keek, Adam Burrows, and Stan Woosley for many useful discussions. The simulations were performed at Minnesota Supercomputer Institute (MSI) and National Energy Research Scientific Computing Center (NERSC).

This project has been supported by the DOE Program for Scientific Discovery through Advanced Computing (SciDAC; grants DOE-FC02-01ER41176, DOE-FC02-06ER41438, and DE-FC02-09ER41618), by the US Department of Energy under grant DE-FG02-87ER40328.

References

1. Timmes (ApJs, 124, 241, 1999).
2. Chen, Heger & Almgren, in preparation.
3. Weaver, Zimmerman, & Woosley (ApJ 225, 1021, 1978).
4. Almgren et al. CASTRO: A New Compressible Astrophysical Solver. I. Hydrodynamics and Self-Gravity, submitted to ApJ.