

Communication, 2.

Recognizing the Message

Distinguishing from natural “signals”:

Expect: Variation with time, narrow band
(small range of freq.)

Crucial → Not random noise

If not random, it is artificial (ETI or Human)

Examples of natural signals that might have been ETI


1. Pulsars (LGM)
2. OH Masers

Both are random noise (no coded information)

Coding the Message


Change the signal with time

1. Amplitude modulation (AM)




Coding the Message

2. Frequency Modulation (FM Radio)



Coding the Message



<http://www.chem.tamu.edu/rgroup/north/FM.html>

Analog vs. Digital

1. Analog - need accurate amplifiers, etc.
to avoid distortion
e.g. radios, tv, records, analog tapes
2. Digital → “digitize” signal
Represent by Base 2 Number

Base 10	Base 2
0	0
1	1
2	10
3	11
4	100
⋮	⋮

Analog vs. Digital

Send one digit at a time so electronics just need to Distinguish 1 from 0

Can use 2 very different voltages


→ Need fast digital electronics

e.g. CD's, DVDs, Computers, Digital Tapes, Digital TV, ...

Decoding the Message

Assume Digital

Repeat to Establish Pattern



Image? 1 dimension (string of bits)



2 dimensions

Rows + columns

Make product of # rows + # of columns
each a prime number


e.g., $23 \times 73 = 1679$ so 23 rows, 73 columns
or vice versa

Semantics

Can we understand the message?

```
0000001010101000000000000101000001010
00000010010001000100010001001011001010101
010101010100010001000000000000000000000
000000000000000000000000000000000000000
110100000000000000000000000000000000000
0000000010101000000000000000000000111110
000000000000000000000000000000000000000
1110001100001100011000100000000000000000
000110100011000110000110001101011110111
011111011111000000000000000000000000000
010000000000000000000000000000000000000
000000000000000000000000000000000000000
000000000000000000000000000000000000000
001100000000000000000000000000000000000
0011000000110000001100001100010000001000
0000001000001101000001100001100011010111
110111101111101111011111000000000000000
000000000010000000110000000000000000000
001100000000000000000000000000000000000
111111100000110000000011111000000000110
000000000000000000000000000000000000001
000000110000000000000000000000000000000
100000000000000000000000000000000000000
011001100000000000000000000000000000000
000011000000000000000000000000000000000
100000010000000000000000000000000000000
000000110000000000000000000000000000000
000000110000000000000000000000000000000
000000000000000000000000000000000000000
011001100000000000000000000000000000000
000000000000000000000000000000000000000
010111010010110110000000000000000000000
111110111000001110000000000000000000000
100000011011001000000000000000000000000
100000001000000000000000000000000000000
000000000000000000000000000000000000000
000100000000000000000000000000000000000
000111111111000000000000000000000000000
000000000011000000000000000000000000000
000000001011000000000000000000000000000
000000001011000000000000000000000000000
001000010100000000000000000000000000000
001000100000000000000000000000000000000
010000100000000000000000000000000000000
000000000000000000000000000000000000000
11101001111000
```

Figure 19.12 The message sent in 1974 from the Arecibo telescope in the direction of the globular cluster M13 consists of 1679 bits of information, either "on" or "off," shown here as 0's and 1's.



Search Strategies

- Basic Problem: where to look?
- Possible Scenarios
 - Powerful, omnidirectional beacons
 - Implies very advanced civilization
 - Seeking to attract attention of new civilizations
 - Nearby, not so advanced, broadcasting to us
 - Unlikely
 - Detect leakage radiation

Leakage Radiation

- Various sources
 - TV, radio, ...
 - Repeatable pattern due to Earth rotation
 - Defense radars
 - Most powerful, but won't repeat

TABLE 20-1
ESTIMATED POWER OUTPUT OF VARIOUS RADIO-PHOTON SOURCES THAT OPERATE AT FREQUENCIES GREATER THAN 20 MHz

Source	Frequency Range (MHz)	Number of Transmitters	Fraction of Time that Transmitters Emit	Per Individual Transmitter		Total Average Power Radiated (watts per hertz of bandwidth) ^a
				Maximum Power Radiated (watts)	Effective Frequency Bandwidth (hertz)	
Citizen-band radios	27	10,000,000	1/100	5	2	200,000
Professional landmobile radios	20-500	100,000	1/10	20	1	200,000
Weather, marine, and air radars	1000-10,000	100,000	1/100	10,000 to 1,000,000	1,000,000	10 to 1000
Defense radars ^b	400	2	1/10	10,000,000,000	0.1	20,000,000,000
FM radio stations	88-108	10,000	1	4000	0.1	400,000,000
TV stations (for photons that carry picture, not sound)	40-850	2000	1	500,000	0.1	10,000,000,000

^aThe last column shows the power radiated *per hertz of bandwidth*. Systems that cover a wider bandwidth (most noticeably, weather, marine, and air radars) will radiate a greater total power over *all* frequencies than this column would suggest. This table, as well as Figures 20-7, 20-8, and 20-9 follow the results of a study made by W. Sullivan III, S. Brown, and C. Wetherill in *Science*, vol. 199, p. 377, 1978.

^bWe have considered only the most powerful defense radars; these dominate the total power output from all such radar systems.







Fig. 3 (left). World television spectrum, show which the most power is radiated (5-Mhz-v primary bands (channels 2 to 6, 7 to 13, and States) are also indicated. A distant extraterrestrial frequency resolution of 5 Mhz would at a spectrum roughly of this shape. Fig. 4 antenna power patterns adopted in the model television frequency bands. The radiation is to the local horizon of each transmitter and directional in azimuth.





The Cosmic Haystack

Frequency

Large frequency range


But narrow channels



Lots of channels

Direction

Large number of directions



Sensitivity

$$S \propto D^{-2} t^{-1/2}$$

want small S

Large telescope

Long time per direction


Strong signals, unknown origin

⇒ Small telescope, short t, cover sky

Weak signals, nearby stars

⇒ Large telescope, longer t, only stars

Cosmic Haystack



INTERSTELLAR RADIO AND TELEVISION MESSAGES






Figure 19.5 Concentrating a signal into a narrower bandpass makes it much easier for the signal to stand out against the background noise that exists at all frequencies.

THE SEARCH FOR EXTRATERRESTRIAL INTELLIGENCE





The Cosmic Haystack

Frequency

Large frequency range


But narrow channels



Lots of channels

Direction

Large number of directions



Sensitivity

$$S \propto D^{-2} t^{-1/2}$$

want small S

Large telescope

Long time per direction

Strong signals, unknown origin

⇒ Small telescope, short t , cover sky


Sky Survey

Weak signals, nearby stars

⇒ Large telescope, longer t , only stars

Targeted Search


Targeted Search vs Sky Survey




Some Searches for ETI

Year	Names	Frequency (MHz)	Telescope size (m)	# of stars
1960	Ozma (Frank Drake)	1420	26	2
1972	Ozma II (Zuckerman & Palmer)	1420	91	602
1985	Meta (Horowitz; Planetary Soc.; Spielberg) [8 million channels]	1420	26	All sky
1992(?) ↑ Oct. 12, 1992	NASA search Discrete source made	{ 1200-3000 + selected ν Up to 25 GHz }	300	244
	All sky Survey		34	800
		1000 - 10,000 + selected ν	34	All Sky
	[10 million channels +?]			
	2 million in 1992 ; ~ 16 million in 1996			

Some Searches



Previous Searches



SERENDIP - SETI@home

- Latest version:
SERENDIP IV
Uses Arecibo telescope
while regular obs.
going on

$\nu = 1420 \text{ MHz}$

$5 \times 10^{-25} \text{ W m}^{-2}$
very sensitive

Data analyzed by screen savers
on millions of PC's SETI@HOME



Report on Project META Megachannel Extra Terrestrial Assay

Horowitz & Sagan, 1993, *Astrophysical Journal*, **415**, 218.

5 years of searching at 1.420 GHz

8×10^6 channels channel width: 0.05 Hz

coverage: 400 kHz

Covered sky 3 times $1.7 \times 10^{-23} \text{ W m}^{-2}$

37 candidate events: narrow-band, apparently not interference

But none repeated


8 signals truly hard to explain as noise

Probably electronic "glitches"

But some tendency to lie in plane of galaxy \Rightarrow extraterrestrial

?

Nothing convincing yet.



BETA

Successor to META 2.5×10^8 channels
0.5 Hz channel width

Covers 1.4 - 1.7 GHz in 8 steps

Sensitivity: 2×10^{-22} W m⁻²

Started 1995, suspended in Spring 1999
 (antenna blew off mount!)
 repairs underway

NASA Search

To ~~begin~~ ^{began} Oct. 12, 1992 ~~ended~~ ^{ended} revived?

Microwave Observing Program (MOP)

Main improvement: frequency coverage

2 parts:

1. All sky survey - JPL - run
Telescopes of modest 34-m diameter
California, Australia, ...
Cover 1 - 10 GHz
 2×10^6 channels 16×10^6 channels (~ 1996)
Channel width: 20 Hz

Coverage: 40 MHz , 320 MHz

right and left circular polarization

Sensitivity: only spend a few sec. per direction

⇒ strong signal

(Arecibo Planetary Radar)

out to 25 ly

Timespan: 6 years to cover sky once

2. Targeted search - Ames - run
 (~ 800 Nearest (< 75 ly) stars like Sun)
 Largest telescopes available:
 Arecibo 300 m (244 stars)
 + Australia, France, ...
 Cover: 1 - 3 GHz
 16×10^6 channels
 Channel width: 1 Hz

Coverage: 10 MHz
 right and left circular polarization
 Sensitivity: $\sim 10^3$ sec. per star
 $\Rightarrow 10^{-27} \text{ W m}^{-2}$
 $P_{\text{trans}} = 10^{-27} \text{ W m}^{-2} \cdot 4\pi d^2(\text{m})$
 $d(\text{m}) \simeq 10^{16} d(\text{ly})$
 $P_{\text{trans}} \simeq 10^6 d^2(\text{ly}) = 1 \text{ M Watt at } 1 \text{ ly}$
 e.g. 100 Mega Watts at $d = 10 \text{ ly}$
 Defense radars to $\sim 1000 \text{ ly}$

HR 5158

EXCERPTS REGARDING SETI 101st Congress of the United States, 2nd Session

From Senate Report 101-474, to accompany H.R. 5158, from the Departments of Veterans Affairs, HUD and Independent Agencies Appropriation Bill, 1991, dated September 16, 1990 (Senator Barbara Mikulski—chair):

Regarding the NASA budget:

... For life sciences, the Committee recommends the following:

-\$25,000,000 from the \$168,000,000 requested for life sciences, to be taken as a general reduction, subject to the normal reprogramming guidelines. None of this reduction is to be taken from the request for the search for extraterrestrial intelligence (SETI) program.

"In recommending the full budget request of \$12,100,000 for the SETI program, the Committee reaffirms its support of the basic scientific merit of this experiment to monitor portions of the radio spectrum as an efficient means of exploring the possibility of the existence of intelligent extraterrestrial life. While this speculative venture stimulates widespread interest and imagination, the Committee's recommendation is based on its assessment of the technical and engineering advances associated with the development of the monitoring devices needed for the project and on the broad educational component of the program. The fundamental character of the SETI program provides unique opportunities to explain principles of such scientific disciplines as biology, astronomy, physics, and chemistry, in addition to exposing students to the development and application of microelectronic technology.

"The Committee has included the full request of \$2,000,000 for the Lifesat project..."

From the Joint House-Senate Conference Report for Veterans Affairs, HUD and Independent Agencies (approved on October 17, 1990):

Regarding the NASA budget:

... \$25,000 from Life Sciences

"The Conferees agree that within the balance of funds available in this section, \$12,100,000 shall be allocated to the Search for Extraterrestrial Intelligence and \$2,000,000 for Lifesat..."

SETI Office/10-90

SEARCH IS DESCRIBED IN CHAP. 9

Sunday, October 10, 1993

Austin American-Statesman A19

Congress may hang up on research of E.T.s

■ Extraterrestrials won't be able to phone home if there's nobody on earth to take the call

By Keay Davidson
New York Times News Service

SAN FRANCISCO — Who killed E.T.?

An effort by the National Aeronautics and Space Administration to detect signals from extraterrestrials has been axed by Congress.

Experts blame everything from its "giggle factor" to poor salesmanship to Congress' unwillingness to cut politically stronger programs.

Hollywood has made big money from fictional extraterrestrials, and they clutter TV shows and grocery-store tabloids.

But NASA's \$104 million attempt to find real aliens — the Mountain View, Calif., High Resolution Microwave Survey — was too costly for a joint congressional committee. It agreed to end the program just one year into its planned 10-year search. The program is popularly known by its previous name, Search for Extraterrestrial Intelligence, or SETI. "I'm pretty depressed," said Pa-

critics accuse Congress of making SETI a sacrificial lamb after failing to kill two programs — the oft-maligned space station and the \$3 billion Advanced Solid Rocket Motor, which Reader's Digest last year called "the unstoppable pork booster." It's based in Yellow Creek, Miss., home to Democratic Rep. Jamie Whitten, who until last year chaired the House Appropriations Committee.

Project staff members took pride in the program's size. "Each space shuttle launch has been estimated to cost as much as \$1 billion. That's a century worth of SETI research," said Seth Shostak of the quasi-private SETI Institute in Mountain View.

But politically, "the SETI people made a fundamental mistake — stupid, stupid, stupid! — in the way they've been lobbying for their programs," said John Pike, a policy expert with the Federation of American Scientists in Washington. "SETI is one of the things that is most readily understood and widely appreciated by the public."

Ralph De Gennaro, a senior budget analyst for Friends of the Earth in Washington, D.C., shed no tears for SETI.

"I'm sick and tired of being told that we can't afford to save this planet but we do have enough money to listen to aliens on other

Project Phoenix

Underway Feb. 2, 1995

SETI Institute (- minus NASA \$\$)

Private Funding (Packard of HP)

+ ...

Relocate to Australia 64 - m telescope

1.2 - 3.0 GHz , 28×10^6 channels

1 Hz channel width

Targeted search sensitivity $\sim 1 \times 10^{-26} \text{ W m}^{-2}$

~ 200 stars like Sun, no binaries, $t \geq 3 \times 10^9$ yr

Within 150 ly observe each for 5 min

(eventually 1000 stars)

Can detect 1 Mega Watt if beamed to us by
similar size telescope

Immediate followup by second telescope

No ETI found in first run (sp 95)

Webpage: <http://www.seti-inst.edu>

Used various other telescopes, including Arecibo

No civilizations found yet.

Amateur Projects

BAMBI (Bob and Mike's Big Investment)

3.7 - 4.2 GHz Sky survey



SETI League project ARGUS

Use Satellite TV Dishes (~ 100) as of 2001

1.4 - 1.7 GHz Channel width: 1 Hz

Sens. $\sim 1 \times 10^{-21} \text{ W m}^{-2}$

Goal is 5000 sites

Aim for continuous sky coverage



Allen Telescope Array (ATA)

Under construction SETI Institute, UC Berkeley
Major telescope dedicated to SETI

Cost $\sim 26 \text{ M } \$$ $\sim 1/2$ provided by Paul Allen,
Nathan Myrrovold (Microsoft)

Hat Creek, California


350 \times 6 m antennas

1 - 10 GHz

Can examine 10^5 stars 3 times over a decade

Will extend targeted search much farther.

Expanding the Search Radius



Comparison of the Allen Telescope Array and Project Phoenix




Figure 2. A representation of the increase in relative detection probability of SETI searches with date. The positive slope of these data is correlated with the technological enhancements that have benefited SETI search systems from one decade to the next.

Websites for SETI

<http://www.seti-inst.edu/> Many Links

<http://www.mc.harvard.edu/seti/> Project BETA

Update on Searches

Article by Jim Tarter, 2001

Annual reviews of Astronomy & Astrophysics, **39**, 511

Appendix Available on WWW

99 SETI projects > 14 ongoing in 2001

Some Optical, most radio

Update on Searches

Notable Ones:

Serendip → SETI@home

META → BETA

NASA → Phoenix

BAMBI, ARGUS (Amateurs)

Allen Telescope Array (Future)

Beyond MOP

VLA Expansion → “ARGUS”

Cyclops

1000 telescopes, each 100-m diameter

Detect 1000 MW transmitter at 1000 μ y

or monitor 1000 stars simultaneously

or detect leakage radiation at 100 μ y

Square Kilometer Array (SKA)

