Laws of Motion

- Aristotle (384-322 B.C.)
 - A body must be continuously pushed by forces in order to keep its motion (idea also adopted by Descartes).
 - *The role of resistance*: the bigger the resistance is, the bigger the applied forces must be.
 - Force = Resistance x Velocity
 - If this law is correct, no vacuum can exist because zero resistance means infinite velocity.
 - Proof of the existence of ether, and disproof of the existence of atoms.

Cartesian Universe

- Descartes (1596-1650)
 - French philosopher, mathematician
 - Cartesian coordinate (x,y,z)
 - Mechanical world of matter and motion which obey natural laws
- The Cartesian universe is infinite and full of matter (no empty space allowed)
 - All motion was explained by direct contact (such as pressures and tensions).
 - Planetary motion was explained by vortical motions of interplanetary matter (or fluid).

Momentum

- No force is necessary to keep motion.
 - A body carries momentum, which keeps its motion.
 - Momentum of a body represents how hard it is to change its motion (speed and/or direction of motion)
 - Momentum = Mass x Velocity
 - Force = Rate of change in momentum
- Angular momentum represents how hard it is to change angular motion of a rotating object.
 - Angular momentum = Distance from the axis of rotation x Momentum
 - Torque = Rate of change in angular momentum
- Important fact: momentum conserves in the absence of force or torque.

Newton's laws of motion

- Isaac Newton (1642-1726)
- 1st law: In the absence of a net force acting upon it, an object moves with constant velocity
 - Momentum (= mass x velocity) of a body is constant in the absence of a net force.
- 2nd law: *Force* = *Rate of change in momentum*
 - Force = Mass x (Rate of change in velocity)
 - Force = Mass x Acceleration
- 3rd law: for any force, there always is an equal and opposite reaction force
 - Total momentum of all bodies is conserved even in the presence of forces.
 - Centrifugal force, weightless astronaut, rocket

Newton's law of gravity

- Gravitational force between two bodies is given by
 - Force = (Gravitational constant) x (Mass of object 1) x (Mass of object 2) divided by squared distance between two objects.
 - $-F = G m M / r^2$
- Discovery of this law was made possible by Kepler's laws.
 - Conflict between Isaac Newton and Robert Hooke:
 "who did it first?"
 - "If I have seen further, it is by standing on the shoulders of giants." -- Letter from Newton to Hooke

Kepler's laws explained by Newton

- Kepler's 1st law: The orbit of each planet about the Sun is an ellipse with the Sun at one focus
 - Natural orbits under the 1/r² force (provided by the Sun) are ellipses, parabolas, and hyperbolas.
- Kepler's 2nd law: As a planet moves around its orbit, it sweeps out equal areas in equal times
 - Conservation of angular momentum (= m r v)
- Kepler's 3^{rd} law: More distant planets orbit the Sun at slower average speeds, obeying the following precise mathematical relation ship: $P^2 = R^3$
 - The 1/r² force from the Sun balances the centrifugal force (Newton's 3rd law), giving this relationship.