


Life

# What is necessary for life?

Most life familiar to us: Eukaryotes

FREE LIVING  
Or Parasites



First appeared  $\sim 1.5 - 2 \times 10^9$  years ago


Requirements: DNA, proteins, lipids, carbohydrates,  
complex structure,  $\sim 10^4 - 10^5$  genes

# Prokaryotes (Bacteria and Archaea)

First appeared


$\sim 3 - 4 \times 10^9$  years ago

FREE LIVING  
Or Parasites




Requirements: DNA, protein, lipids, carbohydrates,  
simpler structure, few thousand genes


# Viruses



Viroids,  
Plasmids




Prions?



The tobacco-mosaic virus is made up of a strand of nucleic acid encased in a rod of one kind of protein.

## TOBACCO MOSAIC VIRUS



# Minimum Requirements for Life

Proteins and Nucleic Acids for simplest  
Or maybe only one.

Lipids and Carbohydrates for any thing  
more complex than a virus.

These are all macromolecules.

# Macromolecules

H, C, N, O  
(S)

Proteins made of amino acids (20)  
Construction and catalysis (enzymes)

H, C, N, O  
(P)

Nucleic acids made of nucleotides

base    sugar    phosphate

Polymers and Monomers

H, C, O

Carbohydrates made of sugars

Energy (food) + structure

[starch]

[cellulose]

H, C, (O)


Lipids (hydrocarbons + carboxyl)

Membranes + Energy

[water-resistant]




# Sugar



Glucose


# Lipids




carboxyl

hydrocarbon

Fatty acid is composed of a hydrocarbon chain with a carboxyl group at one end



# Polysaccharides




# Proteins

Monomers are amino acids

20 kinds


Glycine



Amino  
group


carboxyl  
group

Schematic




Section of Protein


## A Peptide Bond at the Chemical Level




Before




After



Note that a water molecule must be removed




amino acids



protein

A complex protein:


Involved in oxygen use  
Each circle is an amino acid



Stripped down view  
Can you find the  
amino end and the  
carboxyl end?

Note the “heme”,  
containing iron.


Function depends on  
structure, which  
depends on folding,  
which depends on  
order of amino acid  
bases




# Nucleic Acids (DNA, RNA)

Made of sugars, phosphates, bases

Sugar



Schematic




Ribose Sugar  
5 C, 5 O, 10 H


Ribonucleic acid (RNA) uses ribose sugar;  
Deoxyribonucleic acid (DNA) uses deoxyribose sugar



# phosphate



sugars & phosphates linked  
phosphodiester bonds




Segment of side of ladder structure

# Nucleic Acids (cont.)


Bases: Carry Genetic Code

Purines




Adenine

Adenine




Guanine




Equal numbers of C and N

# Pyrimidines




Cytosine

More C than N





Cytosine








Uracil / Thymine

RNA / DNA

## Bases in Nucleic acids: Purines and Pyrimidines






Purines

Pyrimidines


Note Uracil

# Nucleic Acids (cont.)

## Segment of RNA





## Segment of DNA




Note that T  
replaces U in DNA

## At the Chemical Level













# Hydrogen Bonds (weak) connect the bases across the two sides of DNA




# Segment of DNA



The two strands of DNA form a double helix, connected between bases by hydrogen bonds



## Further wrapping to make compact chromosome



# Information Storage

- Nucleic acids store information
- The information specifies proteins
- The information can be replicated
- This allows inheritance

## Base pairing rules

A - T

G - C

- U

⇒ Replication of order  
(reproduction)

Nucleic Acid - Protein

↪ Genetic Code ↩

# Codon

3 base sequence specifies an Amino Acid

# Gene

Sequence of codons specifies a Protein

a gene specifies a protein

e.g.	tobacco mosaic virus	4 genes
	bacteria	$\sim 10^3$ genes
	human cell	$\sim 25,000$ genes (update)

# For mRNA


# Genetic Code

First RNA Base	U	C	A	G	Third RNA BASE
U	Phenylalanine	Serine	Tyrosine	Cysteine	U
	Phenylalanine	Serine	Tyrosine	Cysteine	C
	Leucine	Serine	Stop	Stop	A
	Leucine	Serine	Stop	Tryptophan	G
C	Leucine	Proline	Histidine	Arginine	U
	Leucine	Proline	Histidine	Arginine	C
	Leucine	Proline	Glutamine	Arginine	A
	Leucine	Proline	Glutamine	Arginine	G
A	Isoleucine	Threonine	Asparagine	Serine	U
	Isoleucine	Threonine	Asparagine	Serine	C
	Isoleucine	Threonine	Lysine	Arginine	A
	Start/Methionine	Threonine	Lysine	Arginine	G
G	Valine	Alanine	Aspartic Acid	Glycine	U
	Valine	Alanine	Aspartic Acid	Glycine	C
	Valine	Alanine	Glutamic Acid	Glycine	A
	Valine	Alanine	Glutamic Acid	Glycine	G


Amino Acids





# PROTEIN SYNTHESIS



# Structure of a tRNA



# Translation



# Variations in the Code

## 1. “Wobble” Bases

The third base in a codon can sometimes vary.

tRNA

U

G

mRNA

A or G

C or U

Comparison to genetic code  $\Rightarrow$  no change  
in amino acids

# For mRNA

# Genetic Code

First RNA Base	U	C	A	G	Third RNA BASE
U	Phenylalanine	Serine	Tyrosine	Cysteine	U
	Phenylalanine	Serine	Tyrosine	Cysteine	C
	Leucine	Serine	Stop	Stop	A
	Leucine	Serine	Stop	Tryptophan	G
C	Leucine	Proline	Histidine	Arginine	U
	Leucine	Proline	Histidine	Arginine	C
	Leucine	Proline	Glutamine	Arginine	A
	Leucine	Proline	Glutamine	Arginine	G
A	Isoleucine	Threonine	Asparagine	Serine	U
	Isoleucine	Threonine	Asparagine	Serine	C
	Isoleucine	Threonine	Lysine	Arginine	A
	Start/Methionine	Threonine	Lysine	Arginine	G
G	Valine	Alanine	Aspartic Acid	Glycine	U
	Valine	Alanine	Aspartic Acid	Glycine	C
	Valine	Alanine	Glutamic Acid	Glycine	A
	Valine	Alanine	Glutamic Acid	Glycine	G

Amino Acids

2. Some organisms use slightly different codes, with one or more changes in codon translation.

First seen in mitochondrial DNA.

Now known in some nuclear DNA

The code has evolved since the last common ancestor (But not much).

# Summary

1. Atoms needed: H, C, O, N, small amounts of P (phosphorus), S (sulfur)
2. Two basic molecules needed for life: proteins, nucleic acids
3. Both are polymers - made of simpler monomers. The monomers function as words or letters of alphabet. Information is the key.

## Summary (cont.)

4. Proteins and nucleic acids closely linked at fundamental level. Communicate through genetic code. All organisms have almost the same genetic code. It must have originated very early in evolution of life.
5. In present day organisms, protein synthesis must be directed by nucleic acids, but nucleic acid reading or replication requires enzymes (proteins). Chicken-Egg problem



# Some Movies of Processes

- Animation of transcription (making mRNA)
  - <http://vcell.ndsu.nodak.edu/animations/transcription/movie.htm>
- Animation of translation (making protein)
  - <http://vcell.ndsu.nodak.edu/animations/translation/movie.htm>
- Both from Virtual Cell Animation collection,  
Molecular and Cellular Biology Learning  
Center