


Cosmic Evolution, Part II

Heavy Elements to Molecules

First a review of terminology:



Neutral atom:  
ion:


# Electrons = # protons




e.g.  $\text{C}^{+2}$  Carbon nucleus + 4 (6-2) electrons

# Forces

H atom



H<sub>2</sub>  
Molecule



Attractive

Repulsive




Molecule: Repulsive ~ Attractive

More delicate than atoms,  
can be much more complex

“Bond” is sharing of electrons

Is molecule stable?

Yes, if EM potential energy less than separate atoms






Activation energy lower  $\rightarrow T \sim 100 - 1000 \text{ K}$

(Room Temperature)

# Questions

- Why is room temperature around 300 K?
- How commonly is this temperature found in the Universe?

Conventions:



Maximum # of Bonds:

H      1

O      2

N      3

C      4

Carbon very versatile

→ Complex chemistry


# Interstellar Molecules

Exist as gas (individual molecules)



A few known in 1930's

Many more since 1968 - Radio astronomy


Rotation



Vibration



Radio Telescope




Optical Telescope

# How we detect Interstellar Molecules

Radio Spectroscopy (Mostly  $\lambda \sim 1-3$  mm)

+ Precise knowledge of wavelengths for different molecules





## Appendix 2

### Interstellar Molecules

Species	Name	Species	Name
H <sub>2</sub>	molecular hydrogen	CO <sub>2</sub>	carbon dioxide
C <sub>2</sub>	diatomic carbon	OCS	carbonyl sulfide
CH	methylidyne	SO <sub>2</sub>	sulfur dioxide
CH <sup>+</sup>	methylidyne ion	SiC <sub>2</sub>	silicon dicarbide*
CN	cyanogen	SiCN	
CO	carbon monoxide	AlCN	
CO <sup>+</sup>	carbon monoxide ion	C <sub>2</sub> S	
CS	carbon monosulfide	C <sub>2</sub> O	dicarbon monoxide †
OH	hydroxyl	C <sub>3</sub>	triatomic carbon*
HCl	hydrogen chloride	MgCN	magnesium cyanide*
NH		MgNC	magnesium isocyanide*
NO	nitric oxide	NaCN	sodium cyanide*
NS	nitrogen sulfide		
SiC	silicon carbide*	C <sub>2</sub> H <sub>2</sub>	acetylene
SiO	silicon monoxide	C <sub>3</sub> H	propynylidyne (l and c)
SiS	silicon sulfide	H <sub>2</sub> CO	formaldehyde
SiN	silicon nitride	H <sub>2</sub> CN	
SO	sulfur monoxide	HC <sub>2</sub> N	
PN		NH <sub>3</sub>	ammonia
CP	*	HNCO	isocyanic acid
SO <sup>+</sup>	sulfoxide ion	HOCO <sup>+</sup>	
NaCl	sodium chloride*	HCNH <sup>+</sup>	
AlCl	aluminum chloride*	HNCS	isothiocyanic acid
KCl	potassium chloride*	C <sub>3</sub> N	cycloethynyl
AlF	aluminum fluoride*†	C <sub>3</sub> O	tricarbon monoxide
FeO	iron monoxide	C <sub>3</sub> S	
HF		H <sub>2</sub> CS	thioformaldehyde
SH		H <sub>3</sub> O <sup>+</sup>	hydronium ion
		SiC <sub>3</sub>	
H <sub>3</sub> <sup>+</sup>	protonated hydrogen	C <sub>4</sub> H	butadiynyl
C <sub>2</sub> H	ethynyl	C <sub>3</sub> H <sub>2</sub>	cyclopropenylidene
CH <sub>2</sub>	methylene †	H <sub>2</sub> CCC	propadienylidene
HCN	hydrogen cyanide	HCOOH	formic acid
HNC	hydrogen isocyanide	CH <sub>2</sub> CO	ketene
HCO	formyl	HC <sub>3</sub> N	cyanodiacetylene
HCO <sup>+</sup>	formyl ion	HNC <sub>3</sub>	
HCS <sup>+</sup>	thioformyl ion	CH <sub>2</sub> CN	cyanomethyl
HOC <sup>+</sup>	isoformyl ion †	NH <sub>2</sub> CN	cyanamide
N <sub>2</sub> H <sup>+</sup>	protonated nitrogen	CH <sub>2</sub> NH	methanimine
HNO	nitroxyl	HC <sub>2</sub> NC	
H <sub>2</sub> O	water	CH <sub>4</sub>	methane
H <sub>2</sub> S	hydrogen sulfide		
H <sub>2</sub> N	hydrogen nitride		
N <sub>2</sub> O	nitrous oxide		

Species	Name	Species	Name
H <sub>2</sub> COH <sup>+</sup>	protonated formaldehyde	HC <sub>5</sub> N	cyanodiacetylene
SiH <sub>4</sub>	silane*		
C <sub>4</sub> Si	*	C <sub>7</sub> H	methyl formate
C <sub>5</sub>	pentatomic carbon*	HCOOCH <sub>3</sub>	methyl cyanooacetylene
		CH <sub>3</sub> C <sub>3</sub> N	acetic acid
C <sub>5</sub> H	pentynylidyne	CH <sub>3</sub> COOH	
C <sub>5</sub> N		H <sub>2</sub> C <sub>6</sub>	glycolaldehyde
C <sub>2</sub> H <sub>4</sub>	ethylene*	CH <sub>2</sub> OHCHO	
H <sub>2</sub> CCCC	butatrienylidene		
CH <sub>3</sub> OH	methanol	CH <sub>3</sub> C <sub>4</sub> H	methylidiacetylene
CH <sub>3</sub> CN	methyl cyanide	CH <sub>3</sub> CH <sub>3</sub> O	dimethyl ether
CH <sub>3</sub> NC	methyl isocyanide	CH <sub>3</sub> CH <sub>2</sub> CN	ethyl cyanide
CH <sub>3</sub> SH	methyl mercaptan	CH <sub>3</sub> CH <sub>2</sub> OH	ethanol
NH <sub>2</sub> CHO	formamide	HC <sub>7</sub> N	cyanohexatriyne
HC <sub>3</sub> HO	propynal	C <sub>8</sub> H	
HC <sub>3</sub> NH <sup>+</sup>			
		CH <sub>3</sub> C <sub>4</sub> CN	↑
C <sub>6</sub> H		CH <sub>3</sub> CH <sub>3</sub> CO	acetone
CH <sub>2</sub> CHCN	vinyl cyanide	NH <sub>2</sub> CH <sub>2</sub> COOH	glycine†
CH <sub>3</sub> C <sub>2</sub> H	methylacetylene	CH <sub>2</sub> OHCH <sub>2</sub> OH	ethylene glycol
CH <sub>3</sub> CHO	acetaldehyde		
CH <sub>3</sub> NH <sub>2</sub>	methylamine		
C <sub>2</sub> H <sub>4</sub> O	ethylene oxide	HC <sub>9</sub> N	cyano-octa-tetra-yne
CH <sub>2</sub> CHOH	vinyl alcohol	HC <sub>11</sub> N	cyano-deca-penta-yne

\* Detected in circumstellar envelopes only  
† tentative



Molecular  
Ions

HCO <sup>+</sup>	formyl ion
HCS <sup>+</sup>	thioformyl ion
HOC <sup>+</sup>	isoformyl ion †
N <sub>2</sub> H <sup>+</sup>	protonated nitrogen
HNO	nitroxyl
H <sub>2</sub> O	water
H <sub>2</sub> S	hydrogen sulfide
H <sub>2</sub> N	hydrogen nitride
N <sub>2</sub> O	nitrous oxide


Look at Appendix 2

## Important Examples:


Water



Ammonia



Formaldehyde



Others of Note: CO      Most common after  $\text{H}_2$


HCN,  $\text{HC}_3\text{N}$ , ...  $\text{HC}_{11}\text{N} \rightarrow$  Carbon chains

$\text{CH}_4$       (Methane)

PAHs (Polycyclic aromatic hydrocarbons)

# 3 Lessons

1. Complexity (Up to 13 - atoms) is extraterrestrial  
May be more complex (Hard to detect)  
Glycine ? 1994 so far, not confirmed  
Polycyclic Aromatic Hydrocarbons (PAHs)  
(Infrared evidence)
2. Dominance of Carbon  
Carbon Chemistry not peculiar to Earth
3. Formation & Destruction Analogous to early Earth



Protection by dust grains: scatter and absorb ultraviolet

# Dust particles

Studies of how they scatter and absorb light  
(Ultraviolet → Visible → Infrared)

⇒ Two types, range of sizes up to  $10^{-6}$  m

Carbon

Silicates

PAHs → Graphite

Si + O + Mg, Fe, ...

~ Soot

Both Produced by old stars


# Formation of Interstellar Molecules

## 1. $\text{H}_2$




Must lose the potential energy difference  
before it falls apart ( $\sim 10^{-14}$  s)

Collisions: OK in lab, too slow in space

Emit photon: very slow for  $\text{H}_2$  ( $10^7$  s)



↑  
surface of dust grain



# Formation of Interstellar Molecules

## 2. More complex molecules





Problem is activation energy barrier

$T \sim 10 \text{ K} \ll \text{Barrier}$

Use reactions **without** activation energies

e.g. Molecular ions, like  $\text{HCO}^+$

Cosmic Ray




Energy + simple mol.

→ Reactive mol.


↓

More complex

# Ion - Molecule Reactions



## Molecules on Dust Grains



Stick on grains  
“ice”

Infrared observations show this: as molecules




Vibrate, absorb infrared

e.g.  $\text{H}_2\text{O}$                       absorbs at  $3 \times 10^{-6} \text{ m}$

$\text{CH}_4$                                 absorbs at  $8 \times 10^{-6} \text{ m}$



# Ices on Dust Grains



## Molecules on Dust Grains

Icy “mantles” contain H, O, C, N

Further reactions possible → more complex molecules (e.g. Ethanol)

→ Building blocks of life ?

→ Life ??? Hoyle and Wickramasinghe

New stars and planets form in same regions

# Implications

1. Similar (Carbon-Dominated) Chemistry
2. Direct Role in Origin of Life?
3. Formation + Destruction  
analogous to Early Earth

## Roles of Dust

1. Protection from UV
2. H<sub>2</sub> Formation
3. Freeze-out → Mantles of Ice  
H<sub>2</sub>O, NH<sub>3</sub>, CH<sub>4</sub>, CO<sub>2</sub>, HCOOH, ...  
↑  
Methane

# Star Formation

First factor in Drake Equation: The rate of star formation

# Estimate of Average Star Formation Rate ( $R_*$ ) \*

$$R_* = \frac{\text{\# of stars in galaxy}}{\text{lifetime of galaxy}} = \frac{N_*}{t_{\text{gal}}}$$

$N_*$  : Count them? No

Use Gravity (Newton's Laws)

Sun orbiting center of galaxy at  $250 \text{ km s}^{-1}$  (155 miles per second)

update:  $269 \text{ km s}^{-1}$  reported in Jan. 2009

Kinetic energy =  $\frac{1}{2}$  gravitational potential energy

$$\frac{1}{2} M_{\odot} v^2 = \frac{1}{2} \frac{G M_g M_{\odot}}{R_g} \quad \leftarrow \text{Distance of Sun from center of galaxy}$$

$$\frac{R_g v^2}{G} = M_g$$

## Estimate of Average Star Formation Rate ( $R_*$ )

$$(R_g = 25,000 \text{ ly}) \rightarrow M_g = 1.0 \times 10^{11} M_\odot$$

Update: 28,000 ly gives  $1.4 \times 10^{11} M_\odot$

$$\text{Add stars outside Sun's orbit} \rightarrow M_g \simeq 1.6 \times 10^{11} M_\odot$$

Update:  $2.0 \times 10^{11} M_\odot$

$$N_* \simeq \frac{M_g}{\text{Avg. mass of star}} = \frac{1.6 \times 10^{11}}{0.4} = 4 \times 10^{11} \text{ (} 5 \times 10^{11} \text{)}$$

$$t_{\text{gal}} \simeq 10^{10} \text{ yr (studies of old stars)}$$

$$R_* \simeq \frac{4 \times 10^{11} \text{ stars}}{10^{10}} = 40 \text{ stars per year (5 - 50)}$$

Update: 50 stars per year

## Complicating factors

50 stars per year is an average over history of Milky Way. Current rate is about 5 stars per year. Probably stars formed more rapidly early in history of Milky Way. Any number between 5 and 50 may be correct for our purposes.

Recent work suggests total mass of Milky Way is 3 trillion solar masses (  $3 \times 10^{12} M_{\odot}$  ). This is mostly dark matter outside the orbit of the Sun.

# Star Formation


## Current Star Formation



# Molecular Clouds

- Composition
  - $\text{H}_2$  (93%), He (6%)
  - Dust and other molecules ( $\sim 1\%$  by mass)
    - CO next most common after  $\text{H}_2$ , He
- Temperature about 10 K
- Density (particles per cubic cm)
  - $\sim 100 \text{ cm}^{-3}$  to  $10^6 \text{ cm}^{-3}$
  - Air has about  $10^{19} \text{ cm}^{-3}$
  - Water about  $3 \times 10^{22} \text{ cm}^{-3}$
- Size 1-300 ly
- Mass 1 to  $10^6 M_{\text{sun}}$

# A Small Molecular Cloud



ESO PR Photo 20a/99 (30 April 1999)

The "Black Cloud" B68  
(VLT ANTU + FORS1)

© European Southern Observatory




# Current Star Formation

- Occurs in gas with heavy elements
  - Molecules and dust keep gas cool
  - Radiate energy released by collapse
  - Stars of lower mass can form
  - Mass needed for collapse increases with  $T$
- Star formation is ongoing in our Galaxy
  - Massive stars are short-lived
  - Star formation observed in infrared

## The Launch of The Spitzer Space Telescope



Spitzer Space Telescope Launched Aug. 2003, expect a 5 yr life.



C. Lachata

# Visible to Infrared Views












RCW 49

234

## Artist's Conception




Features:  
Dusty envelope  
Rotation  
Disk  
Bipolar outflow

R. Hurt, SSC

# The Protostar

- Evolution of the collapsing gas cloud
  - At first, collapsing gas stays cool
  - Dust, gas emit photons, remove energy
  - At  $n \sim 10^{11} \text{ cm}^{-3}$ , photons trapped
  - Gas heats up, dust destroyed, pressure rises
  - Core stops collapsing
  - The outer parts still falling in, adding mass
  - Core shrinks slowly, heats up
  - Fusion begins at  $T \sim 10^7 \text{ K}$
  - Protostar becomes a main-sequence star

# The Disk



The Star (AU Mic) is blocked in a coronagraph. Allows you to see disk. Dust in disk is heated by star and emits in infrared.

# Angular Momentum

- Measure of tendency to rotate
  - $J = mvr$
- Angular momentum is conserved
  - $J = \text{constant}$
  - As gas contracts ( $r$  smaller),  $v$  increases
  - Faster rotation resists collapse
  - Gas settles into rotating disk
  - Protostar adds mass through the disk

# The Wind

- Accretion from disk will spin up the star
  - Star would break apart if spins too fast
- Angular momentum must be carried off
- The star-disk interaction creates a wind
- The wind carries mass to large distances
  - $J = mvr$ , small amount of  $m$  at very large  $r$
  - Allows star to avoid rotating too fast
- Wind turns into bipolar jet
  - Sweeps out cavity

# The Bipolar Jet



**Embedded Outflow in HH 46/47**


NASA / JPL-Caltech / A. Noriega-Crespo (SSC/Caltech)

**Spitzer Space Telescope • IRAC**

Infrared visible light (055)


ssc2003-06f

# Studying the Disk



Robert Hurt, SSC





## Ices in a Protoplanetary Disc

NASA / JPL-Caltech / K. Pontoppidan (Leiden Observatory)

Spitzer Space Telescope • IRS

ESO • VLT-ISAAC

ssc2004-20c