


## Planet Detection

Estimating  $f_p$

## Can We See Them?

- Not easily
  - Best cases were reported in late 2008
  - Will see these later...
- Problem is separating planet light from star light
  - Star is  $10^9$  times brighter in visible light
  - “Only”  $10^6$  times brighter in infrared


## Planet is Much Fainter than Star




## Indirect Detection

### Wobbling star


Detect effect of planet on star (both orbit around center of mass)



Large planet will make a star “wobble”



## Star and Planet Orbit Center of Mass



## The Astrometric Technique

Measure stellar position (angle) accurately - see wobble compared to more distant stars

How far does the star wobble?  
Center of mass

$$M_* \quad M_{pl}$$

$$R_* = \frac{M_{pl} r}{M_*}$$


We measure angle; for small angles,

$$\theta = \frac{R_*}{D} \text{ in radians}$$

$$\text{so } \theta = \frac{M_{pl} r}{M_* D} \quad \begin{matrix} \text{Big planet, big orbit} \\ \text{small star, close to sun} \end{matrix}$$

$$\text{Current limit: } 1 \text{ mas} = 10^{-3} \text{ arcsec} = 2.8 \times 10^{-6} \text{ degrees} = 4.9 \times 10^{-8} \text{ radians}$$

$$\text{e.g. } M_{pl} = M_{Jupiter}, M_* = M_{\odot}, D = 15 \text{ ly} \Rightarrow \theta = 1 \text{ mas}$$



The Sun as viewed from 10 pc (~30 ly)


30 ly

Planet	$M_p$ ( $M_J$ )	$R$ (AU)	$P$ (years)	$V_{\star}$ ( $\text{m s}^{-1}$ )	$\theta$ at 10 pc (mas)
Mercury	1.74E-4	0.387	0.241	0.008	6.4E-6
Venus	2.56E-3	0.723	0.615	0.086	1.8E-4
Earth	3.15E-3	1.000	1.000	0.089	3.0E-4
Mars	3.38E-4	1.524	1.881	0.008	4.9E-5
Jupiter	1.0	5.203	11.86	12.4	0.497
Saturn	0.299	9.54	29.46	2.75	0.273
Uranus	0.046	19.18	84.01	0.297	0.084
Neptune	0.054	30.06	164.8	0.281	0.156
Pluto	6.3E-6	39.44	247.7	3E-5	2.4E-5

## The Spectroscopic Method

- Relies on Doppler Effect
- Motion of star towards and away from us
- Most planets around other stars found by this method so far

## The Doppler Shift



## The Spectroscopic Technique

Measure velocity, not position, of star

Use spectrometer to get Doppler Shift of spectral line


light

shift

$\lambda$

$$\text{Shift} \propto V_{\star} \propto \frac{M_p}{M_{\star}^{1/2}} r^{1/2}$$

Big planet, small orbit  
Small star  
Distance doesn't matter (except for brightness)  
Edge - On



Motion of the Sun caused by Jupiter, ...

30 ly

Planet	$M_p$ ( $M_J$ )	R (AU)	P (years)	$V_*$ ( $m s^{-1}$ )	$\Theta$ at 10 pc (mas)
Mercury	1.74E-4	0.387	0.241	0.008	6.4E-6
Venus	2.56E-3	0.723	0.615	0.086	1.8E-4
Earth	3.15E-3	1.000	1.000	0.089	3.0E-4
Mars	3.38E-4	1.524	1.881	0.008	4.9E-5
Jupiter	1.0	5.203	11.86	12.4	0.497
Saturn	0.299	9.54	29.46	2.75	0.273
Uranus	0.046	19.18	84.01	0.297	0.084
Neptune	0.054	30.06	164.8	0.281	0.156
Pluto	6.3E-6	39.44	247.7	3E-5	2.4E-5

### What We Can Learn

1. There is a planet  
(If not a mistake)
2. The orbital period (P)  
(The time for pattern to repeat)
3. The orbital radius  
 $r^3 \propto M_* P^2$   
(Kepler's Third Law)
4. Lower limit to planet mass ( $M_{planet}$ )


Conservation of momentum

$$M_{pl} > \frac{M_* V_* P}{2\pi r}$$

= if we see orbit edge-on  
> if tilted

### Comparison of Search Methods

Advantages	Astrometric	Spectroscopic
	Big Planet	Big Planet
	Big Orbit	Small Orbit
	Small Star	Small Star
	Nearby Star	--
		Edge-on Orbit



### Other Methods

Transits: Planet passes in front of a star

US

Light from star

Only about 0.5% of stars with planets will line up

planet

star

transit

Time

First planet found with this method in January 2003; 35 detected as of January 2008, 55 by January 2009

Microlensing: Light from more distant star is focused by gravity of nearer star passing in front

US

Fortuitous alignment  $\Rightarrow$  brightens

nearer star


distant star

light from distant star

planet

8 planets found this way as of January 2009

### Artist's conception of Transit of HD209458



Copyright Lynette Cook  
used with permission

<http://www.extrasolar.spaceart.org>


To simulate, try [http://planetquest.jpl.nasa.gov/gallery/gallery\\_index.cfm](http://planetquest.jpl.nasa.gov/gallery/gallery_index.cfm)

### Timing

- Delays or advances in periodic signals
  - Pulses from pulsar
    - First planets found that way in 1992
    - Not suitable for life!
  - Oscillations in white dwarfs
    - First found this way in 2007 by grad student at UT


### Planets from the Transit Method

OGLE-TR-10 P=3.10140 (days)




Light curve

OGLE-TR-10



Star field, shows star

## Planet Detected by Microlensing



OGLE 2005-BLG-235Lb, announced 1/25/06

<http://www.eso.org/outreach/press-rel/pr-2006/pr-03-06.html>

## Current Statistics (Jan. 2011)

- Based on Extrasolar Planets Encyclopedia – <http://exoplanet.eu/>
- 518 Planets
- 54 stars with multiple planets
- Most planets in one system is 5 (55 Cancri)
- Least massive
  - $M = 0.01 M_{\text{Jup}} = 3.0 M_{\text{earth}}$  (MOA-2007-B)
  - $M = 1.8 M_{\text{earth}}$  (Gl 581e, but in question)

## Implications of Planets around other stars

Planets more massive than Jupiter can form around stars like the Sun.

Large Planets can form much closer to a star than Jupiter (or move there)

Does this mean we are unusual and our ideas about other planetary systems are just “solar system chauvinism”?


Not necessarily.

The ones found so far are the “easy” ones. (Big planets close to a star)

Now there are many more with lower masses than higher masses.

Too early to say that we are unusual.

## Number of planets for different masses



### Estimating $f_p$

- Maximum?  $f_p \sim 1$ 
  - All young stars may have disks
- Binaries?
  - Can have disks, but planet formation?
  - Even if form planets, orbits may not be stable
  - If reject binaries,  $f_p < 0.3$

### Estimating $f_p$

- Minimum?
  - Based on success rate of searches ( $n_{\text{found}}/n_{\text{searched}}$ )
  - Extrapolate trends to finding
    - Smaller planets, larger orbits, ...
  - Estimates range from 0.3 to 0.7
    - Note much larger than 0.02 given in book
- Allowed range:  $f_p = 0.3$  to  $1.0$ 
  - Explain your choice!
  - Include/exclude binaries?



### Ongoing Missions and Future Prospects

#### Transits

CoRoT Dec. 2006-present

Has reported numerous new planets

Kepler (Launched March 2009)

Monitor 100,000 stars for 4 years

"Hundreds of Terrestrial Planets"

So far, 11 announced, 700 candidates


Most notable: Kepler 10b, first rocky planet, 1.4 times size of Earth

#### Astrometric Method

GAIA Launch planned for 2012

$M_J$  Planets out to 600 ly; predicted to detect 15,000 planets


## Direct Detection NOW!



Three planets detected with ground-based telescopes working in the near-infrared. They used adaptive optics and other techniques to block the direct starlight and obtain very good spatial resolution.

Three planets at 24, 38, and 68 AU. Evidence for orbiting the star. Best guess masses 5-14  $M_{\text{Jupiter}}$ . Reported in Science, 2008, 322, 1348 by Marois et al.

## And, in the same issue...



A planet just inside the dust ring around the star Fomalhaut. A coronagraph was used to block the star light and the Hubble Space Telescope avoids seeing. The faint spot in the box moves across the sky with Fomalhaut (proper motion), but with a slight shift due to its orbital motion (see inset). Announced by Kalas et al. Science, 2008, 322, 1345.

Planet about 3  $M_{\text{Jupiter}}$  and 119 AU from star. Keeps inner edge of dust ring sharp. Light may actually be reflected off a circumplanetary disk.

## Direct Detection in Future

- Terrestrial Planet Finder (TPF)
  - Under study, not planned for this decade.
  - TPF-C Visible light coronagraph
  - TPF-I Infrared interferometer
- Goal is to detect earth-mass planets
- And to see what gases in atmosphere
  - Suitable for life?
- [http://planetquest.jpl.nasa.gov/TPF/tpf\\_index.html](http://planetquest.jpl.nasa.gov/TPF/tpf_index.html)

## TPF Concepts


TPF-I Infrared Interferometer (2020)



TPF-C Visible light coronagraph (2014)



## Spectroscopy of atmosphere



## Planet Detection Methods

Michael Perryman, Rep. Prog. Phys., 2000, 63, 1209 (updated November 2004)  
[corrections or suggestions please to michael.perryman@esa.int]

