


Origin of Planets

Our Solar System as Example

- We know far more about our solar system than about any other
- It does have (at least) one planet suitable for life
- Start with facts about the solar system
- Then discuss theories of planet formation

General Properties of the Solar System

- Dynamical Regularities
 - Planet orbits in plane, nearly circular
 - Planets orbit sun in same direction (CCW as seen from North Pole)
 - Rotation axes perpendicular to orbit plane
 - Uranus is the exception
 - Planets contain 98% of the angular momentum
 - The Sun contains 99.9% of the mass



Spacing and Composition

- Spacing increases with distance
 - Evenly spaced if plotted logarithmically
 - Missing planet in asteroid belt
- Composition varies with distance
 - Inner 4 are “terrestrial”
 - Small, rocky, thin atmospheres
 - Outer 4 are “gas giants”
 - Gaseous, large, mostly atmosphere

$$\frac{d_2}{d_1} \approx 1.5 \cdot 2$$

The Solar System

Distance from Sun .1 1 10 100 AU (Log scale)

M (M_⊙) 0.06 1 <.001 95 17

Terrestrial { Gas Giants | Icy dwarfs

Asteroids

Composition (%)	Enhancement to get ⊙ abundance		
	Rocky	“Icy”	Gaseous
Terrestrial	100	<1	0
Jupiter	6	~13	~81
Saturn	21	~45	~34
Uranus	~28	~62	~10
Neptune	~28	~62	~10
Comets	~31	~69	~0





Rocky - iron, silicates, ...
 “Icy” - at time of formation (H₂O, NH₃, CH₄, ...)
 Gaseous - H, He


What is a Planet? I. Small end...

- Pluto much smaller than others (0.002 M_{earth})
- Other, similar objects found in Kuiper Belt
 - Including one similar to Pluto (Eris)
 - First named Xena, renamed Eris, goddess of discord, has a moon, Dysnomia, goddess of lawlessness...
- IAU voted in 2006
 - 1. Create a new category of dwarf planet
 - 2. Demote Pluto to a dwarf planet


Theories of Planet Formation

- All start with rotating disk
 - Mass 0.01M_⊙ or more
 - Sum of planet masses 0.001 M_⊙
 - Consistent with observed disk masses
 - Temperature and Density decrease with distance from forming star
 - Dust plays crucial role






Formation of Gas Giants (Jupiter, Saturn)



Predictions from Models


- Formation in rotating disk with icy dust can explain many facts about our solar system
- If we can generalize, expect planetary systems common
- Expect (?) about 10 planets, terrestrial planets in close, giant planets farther out, spaced roughly logarithmically
- May still be typical, but not universal...
- Big planets may clear a gap in disk

Theory Predicts Forming Planets Clear a Gap



Can we observe such gaps?

Detecting Disk Gaps




Disk with little dust

Disk with little hot dust, more cool dust

Disk with almost no hot dust, but a lot of cool dust


Star emission peaks at visible/near-infrared wavelengths. Dust in disks emits in infrared (an excess). Near-infrared if hot (close to star), mid-infrared if cooler. Some excesses start only at long wavelengths but are substantial. These fit models with gaps.

A Case Study LkHα330




Some excess at short λ , but much more beyond 20 μm . Blue line has no gap, red has gap. Implies large gap; models predict about 40 AU radius. Submm interferometer should show ring. J. Brown et al. 2007

Brown et al. 2007b



Detection of planet within disk



A planet just inside the dust ring around the star Fomalhaut. A coronagraph was used to block the star light and the Hubble Space Telescope avoids seeing. The faint spot in the box moves across the sky with Fomalhaut (proper motion), but with a slight shift due to its orbital motion (see inset). Announced by Kalas et al. Science, 2008, 322, 1345.

Planet about $3 M_{\text{Jupiter}}$ and 119 AU from star. Keeps inner edge of dust ring sharp. Light may actually be reflected off a circumplanetary disk.


Issues for Planet Formation

- The time to build up the giant planets from dust particles is long in theories
 - Gas has to last that long to make gas giants
- How long do dust disks last?
 - How long does the gas last?
- Are there faster ways to make planets?
- What about planet building for binary stars?


Time Available to form planets

- The disks around young stars can form planets
- How long do the disks last?
 - Sets limit on time to form planets
 - Half of disks have little dust left by 2 Myr
 - Most gone by 3 to 5 Myr
 - No evidence that gas stays longer
 - Some “debris” around older stars
 - May be evidence of planet building

Disks versus Age of Star Evidence for Collisions



Formation of Gas Giants (Jupiter, Saturn)



Brown Dwarfs

- Stars range from 0.07 to $\sim 100 M_{\text{sun}}$
- Jupiter is about $0.001 M_{\text{sun}}$
- Brown dwarfs between stars and planets
 - Dividing line is somewhat arbitrary
 - Usual choice is $13 M_{\text{Jupiter}}$
 - Brown dwarfs rarely seen as companions to stars
 - But “free-floaters” as common as stars
 - Many young BDs have disks
 - Planets around BDs?

What is a Planet? II. High end...

- Brown dwarfs now found to very low masses
 - Some clearly less than $13 M_{\text{jupiter}}$
 - Can't even fuse deuterium
 - Some people call these planets
 - Some are less massive than known planets
 - Usual definition: planets orbit stars
 - Some brown dwarfs may have "planets"
- Nature does not respect our human desire for neat categories!

Binary Stars

- About 2/3 of all stars are in binaries
 - Most common separation is 10-100 AU
- Can binary stars have disks?
 - Yes, but binary tends to clear a gap
 - Disks well inside binary orbit
 - Or well outside binary orbit

Other Active Issues


- Other planetary systems are quite different
 - Big planets in close
 - But this is probably due to selection effect
- Locations may differ with mass of star
 - Ices survive closer to lower mass star
 - May get ice giants in close
 - Also planets may migrate inwards
 - May prevent formation of terrestrial planets

Formation of Earth


- Almost entirely rocky material (iron, silicates)
- Radioactive elements heat interior
 - Were produced in supernovae explosions
- Interior becomes molten, iron sinks to core
 - Releases gravitational potential energy
 - Interior even hotter
 - Differentiated planet
- Collision forms Earth-Moon system
- Earth acquires atmosphere
 - Outgassing and delivery by comets

Radioactive Heating

Radioactive nuclei decay (release of nuclear potential energy)



Results in layered Earth (like a soft-boiled egg)



- Egg
- Yolk { 1. Inner solid core } T up to 7000 K, iron, nickel, ...
 - 3. Outer liquid core
 - White { 4. Lower mantle - iron rich silicates, solid
 - 5. Asthenosphere (upper mantle) pliable
 - Shell 6. Lithosphere - rigid silicates (crust)
- Lithosphere can "float" on asthenosphere
Leads to Continental Drift, Earthquakes, Volcanos

Continental Drift Reconstructed



150 Myr Reconstruction
Shows motion of continental plates over last 150 Myr.
Red and green dots show locations of ocean drilling.

<http://www.ods.de/ods/index.html>

Formation of Earth and Moon

$$\frac{M_{\text{Moon}}}{M_{\text{Earth}}} \quad \text{Larger than all other planets}$$

Most terrestrial planets have no moons
(Marian moons are captured asteroids)

Moon most likely resulted from giant impact



Earth gets more iron $\rho_{\oplus} = 5.5 \text{ g cm}^{-3}$

Moon mostly silicate $\rho_{\text{Moon}} = 3.3 \text{ g cm}^{-3}$

Temperature was very high after impact (10,000 - 60,000 K)

Any icy material left?


Origin of Atmosphere

Certain "Noble" gases (e.g. Neon) are more rare in Earth atmosphere than in solar nebula. \Rightarrow Atmosphere not collected from gas

Reason: Earth is small \Rightarrow gravity is weak

Temperature in solar nebula is high - atoms moving fast, harder to hold

Outgassing: "Icy" material vaporized by high temperatures



No O₂ on early Earth; No ozone (O₃), so no protection from ultraviolet light

Alternative: Icy materials brought by comets.

Summary

- Planet formation theory suggests planetary systems should be common
- We see evidence of planet building in disks around young stars
- Binary stars with separations around 10 AU may prevent planet formation
- Consider these facts in estimating f_p
- Inner, terrestrial planets of sufficient size will differentiate, may get atmosphere, oceans
- Earth may be unusual (big Moon by collision)