Communication, 2.

Search Strategies


- · Basic Problem: where to look?
- Possible Scenarios
 - Powerful, omnidirectional beacons
 - Implies very advanced civilization
 - Seeking to attract attention of new civilizations
 - Nearby, not so advanced, broadcasting to us
 - Unlikely
 - Detect leakage radiation


Leakage Radiation


- · Various sources
 - TV, radio, ...
 - Repeatable pattern due to Earth rotation
 - Switch to digital TV in June 2009
 - Some changes, but similar frequencies used
 - Defense radars
 - Most powerful, but won't repeat


	Frequency Range (MHz)	Number of Transmitters	Fraction of Time that Transmitters Emit	Per Individual Transmitter		
Source				Maximum Power Radiated (watts)	Effective Frequency Bandwidth (hertz)	Total Average Power Radiated (watts per hertz of bandwidth)*
Citizen-band radios	. 27	10,000,000	1/100	5	2	200,000
Professional landmobile radios	20-500	100,000	1/10	20	1	200,000
Weather, marine, and air radars	1000-10,000	100,000	1/100	10,000 to 1,000,000	1,000,000	10 to 1000
Defense radars ^b	400	2	1/10	10,000,000,000	0.1	20,000,000,000
FM radio stations	88-108	10,000	1 .	4000	0.1	400,000,000
TV stations (for photons that carry picture, not sound)	40-850	2000	1	500.000	0.1	10,000,000,000


"The last column shows the power radiated per here; of bandwidth. Systems that cover a wider handwidth (most noticeably, weather, marine, an irradars) will radiate a greater footh power over all frequencies than this column would suggest. This table, as well as Figures 20-7, 20-8, and 20 9 follow the results of a study made by W. Sullivan III, S. Brown, and C. Wetherill in Science, vol. 199, p. 377, 1978.
































<u>Year</u>	<u>Names</u>	Frequency (MHz)	Telescope size (m)	# of stars
1960	Ozma (Frank Drake)	1420	26	2
1972	Ozma II (Zuckerman & Palmer)	1420	91	602
1985	Meta (Horowitz; Planetary Soc.; Spielberg) [8 million channels]	1420	26	All sky
1992 ↑	NASA search Discrete source made	1200-3000 + selected v Up to 25 GHz	300	244
] Oct. 12, 199		Up to 25 GHz	→ 34	800
	All sky Survey	1000 - 10,000 + selected v	34	All Sky
	[10 million channels +?]			







SERENDIP - SETI@home

 Latest version: SERENDIP IV Uses ARECIBO telescope while regular obs. going on

v = 1420 MHz

 $5 \times 10^{-25} \text{ W m}^{-2}$ very sensitive

Data analyzed by screen savers on millions of PC's SETI@HOME



Report on Project META Megachannel Extra Terrestrial Assay

Horowitz & Sagan, 1993, Astrophysical Journal, 415, 218.

5 years of searching at 1.420 GHz

 8×10^6 channels channel width: 0.05 Hz

400 kHz coverage:

Covered sky 3 times 1.7×10^{-23} W m⁻²

37 candidate events: narrow-band, apparently not interference


But none repeated

8 signals truly hard to explain as noise

Probably electronic "glitches"

But some tendency to lie in plane of galaxy ⇒ extraterrestrial

Nothing convincing.



BETA

Successor to META

 2.5×10^8 channels

0.5 Hz channel width

Covers 1.4 - 1.7 GHz in 8 steps

Sensitivity: $2 \times 10^{-22} \text{ W m}^{-2}$

Started 1995, stopped in Spring 1999 Antenna blew off mount, since dismantled. NASA Search

began ended revived?
To begin Oct. 12, 1992
ing Proces Microwave Observing Program (MOP) Main improvement: frequency coverage 2 parts:

All sky survey - JPL - run

Telescopes of modest 34-m diameter

California, Australia, ...

Cover 1 - 10 GHz

 2×10^6 channels 16×10^6 channels (~ 1996)

Channel width: 20 Hz

Coverage: 40 MHz, 320 MHz

right and left circular polarization

Sensitivity: only spend a few sec. per direction

 \Rightarrow strong signal

(Arecibo Planetary Radar)

out to 25 *l*y

Timespan: 6 years to cover sky once

Targeted search - Ames - run
 (~800 Nearest (<75 ly) stars like Sun)
 Largest telescopes available:

Arecibo 300 m (244 stars)

+ Australia, France, ... Cover: 1 - 3 GHz 16×10^6 channels Channel width: 1 Hz

Coverage: 10 MHz

right and left circular polarization

Sensitivity: ~ 10³ sec. per star

 \Rightarrow 10⁻²⁷ W m⁻²

 $P_{trans} = 10^{-27} \text{ W m}^{-2} \cdot 4\pi \ d^2(m)$

 $d(m) \simeq 10^{16} d(\ell y)$

 $P_{trans} \simeq 10^6 \ d^2(\ell y) = 1 \ M$ Watt at 1 ℓy e.g. 100 Mega Watts at d = 10 ℓy Defense radars to \sim 1000 ℓy

EXCERPTS REGARDING SETI

101x Congress of the United States, 2nd Session

Elemantic States (1997) And Congress of the United States, 2nd Session

Elemantic States (1997) And Congress of Second Segments (1997) And Second Segments (1997)

(Elemantic States (1997) And Second Segments (1997) And Second Segments (1997)

Engaging in NASA bedges:

- 1. For United States, the Considerer recommends the following:
- 4.555,000 from the 1000,000 from the states in the following:
- 4.555,000 from the 1000,000 from 1000,000



Project Phoenix 1998-2004

SETI Institute (- minus NASA \$\$)
Private Funding (Packard of HP)
Relocated to Australia 64 - m telescope
Used various other telescopes, including Arecibo
1.2 - 3.0 GHz, 28 × 10⁶ channels, 1 Hz channel width
Targeted search: 850 nearby stars within 240 &
Sensitivity ~ 1 × 10⁻²⁶ W m⁻²
Could detect 1 Mega Watt if beamed to us by similar size

telescope
Used a second telescope to discriminate against

interference No civilizations found

Amateur Projects

BAMBI (Bob and Mike's Big Investment) 3.7 - 4.2 GHz Sky survey 1997-1999, but may still be going



SETI League project ARGUS
Use Satellite TV Dishes
1.4 - 1.7 GHz Channel width: 1 Hz
Sens. ~ 1 × 10⁻²¹ W m⁻²
About 100 sites in 2000

About 100 sites in 2000 Aim for continuous sky coverage Current status?



Allen Telescope Array (ATA)

SETI Institute, UC Berkeley
Major telescope dedicated to SETI
Partially constructed, some operations (2006)
Cost ~ 26 M \$ ~ 1/2 provided by Paul Allen,
Nathan Myrvold (Microsoft)
Hat Creek, California 350 × 6 m antennas
1 - 10 GHz

Began operation with 42 telescopes in Oct. 2007


Goals for Allen Telescope Array

Survey 1,000,000 stars for non-natural extraterrestrial signals with enough sensitivity to detect the equivalent power of the Arecibo radar out to 1000 light-years within the frequency range of 1 to 10 GHz

Survey the 4×10^{10} stars of the inner Galactic Plane in the "water hole" frequency range from 1420 MHz to 1720 MHz for very powerful, non-natural transmitters

They need funding for more antennas.

Expanding the Search Radius



Comparison of the Allen Telescope Array and Project Phoenix

Searches with Visible Light

For pulsed signals, visible light from lasers. Some advantages. Can concentrate light in narrow band, short pulses to distinguish from star light. They have to be beamed toward us.

Optical SETI at Harvard: 1.5-m telescope 4 x 10^{-9} W m $^{-2}$ in nanosecond pulses Plan to observe 13000 stars. Also a northern sky survey.

Optical SETI in California 1-m telescope Multiple detectors to avoid false signals.

Websites for SETI

SETI Institute list of searches

Many Links and lists of projects

Some out of date

Future Dreams

- Square Kilometer Array (SKA)
 - Use many smaller units
 - Total area about 1 square km
 - Similar to, but beyond, "Argus" in Contact
 - Probably will be three separate arrays
 - To cover full range: 100 MHz to 50 GHz

