April 11, 2011

Exam 4, Friday, April 15. Review sheet posted.

Review session Thursday, 5 - 6 PM Room TBD.

Reading: Chapter 9, Sections 9.5.2, 9.6.1, 9.6.2. 9.7, 9.8

Chapter 10, Sections 10.1-10.6, 10.9

Chapter 11, Sections 11.1 - 11.5

Astronomy in the news: wild fire near McDonald Observatory, 15 homes, 8 outbuildings destroyed in Fort Davis, power, internet, telescopes down at the Observatory. Observatory played host Saturday night to evacuees from Ft. Davis, state parks, tourists.

Pic of the day: Venus rising, Canyon Lands National Park, Utah.



Goal:

To understand how we have discovered supermassive black holes and how they affect galaxy formation and evolution.

Supermassive Black Holes

Long suspected in quasars, active galactic nuclei: huge power from small volume, billion solar mass black hole could do it.

More recently, proof that many (even most! John Kormendy, UT) ordinary galaxies also have a supermassive black hole in their centers (dead quasar).

Again, do not yet see a "dark spot," but use Kepler's Laws, motion of many stars, gas ⇒ orbital period, separation

3.7 million M_☉ black hole in our Galaxy [UCLA link - movie]

Center of Milky Way Galaxy in direction of constellation Sagittarius – (find Sagittarius for sky watch)

Up to billion M_{\odot} black holes in quasars.

Jet from billion M_{\odot} black hole in center of M87, large elliptical galaxy in the Virgo cluster (find Virgo!)

Surprising discovery:


It was long thought that supermassive black holes were somewhat incidental to galaxies,


Formed of matter that somehow drained into the center of the galaxy, so galaxy could have large mass or small mass black hole depending on circumstances.

Recent work by Karl Gebhardt (UT) and others has shown that even stars so far from the center that they cannot possibly feel the gravity of the black hole *now* are moving in such a way that *the larger the mass black hole, the higher the speed of the stars!*

Andromeda galaxy M31

Find Andromeda for sky watch





Mass of Central Bulge of Galaxy


The implication is that the mass of the galaxy (at least the inner portions, the Bulge) is always close to 800 times the mass of the black hole.

This means that the formation of the black hole is somehow intimately connected with the formation and structure of the whole galaxy.

Galaxies "know" how big a black hole to make.

Mechanism uncertain: Does the galaxy control the black hole or the black hole somehow control the galaxy?


Most popular current idea: energy from accretion of matter into disk around black hole feeds back to the surrounding galaxy, blowing excess galaxy gas away when galaxies are young and growing. Colliding black holes in 3C75, feed energy back into the stars and gas of the colliding galaxies.



One Minute Exam

How can we discover a stellar mass black hole that has no accretion disk around it?






Look for jets

We can't

One Minute Exam

What is the relation between the mass of a supermassive black hole and the galaxy in which it resides?



The larger the mass of the galaxy, the smaller the mass of the black hole

The larger the mass of the galaxy, the larger the mass of the black hole


The larger the radius of the galaxy, the larger the mass of the black hole

Goal:


To understand the nature of cosmic gamma-ray bursts, how they may represent the birth of black holes or magnetars, and how they are connected with Type Ic supernovae.

Gamma-Ray Bursts (Chapter 11)

Cosmic explosions, flashes of gamma-rays lasting about 30 seconds, detected by satellites.



Seen across the Universe.



Energy is expelled in narrow jets. Energy comparable to that of supernovae, but all in gamma-rays, with later afterglow in X-ray, radio and optical radiation.

Birth of a black hole?



Swift satellite




Gamma-Ray Bursts unite stars and cosmology

Mystery since late 60's - satellites to monitor space nuclear test ban treaty, avoid confusion between astronomical effects, and bombs

Flare of γ -rays lasts ~ 30 sec

Never Repeat - for 30 years, no optical counterpart, Can't focus gamma-rays. Did not know which of millions of stars to look at.




Did not know the distance: guesses ranged from within the Solar system to cosmologically distant

Goal:

To understand what a gamma-ray burst "afterglow" is and why it is so important.

Revolution in 1997: 1st detection of "afterglow" - optical, radio, X-ray, fading light



Position localized - could bring full armament of modern astronomy to bear on the fading radiation.

⇒Found bursts were in distant galaxies - all at huge, cosmological distances, billions of light years away.

⇒ Very bright to shine that far

January 23, 1999 optical flash associated with the gamma-ray burst itself (need to discover, swivel telescope, look in 30 seconds!)

9th magnitude - human limit 6th magnitude, could almost see with naked eye, could have seen with good binoculars, but half way across the Universe!

March 19, 2008, "naked-eye" GRB 080319B discovered by Swift satellite had a peak apparent magnitude of 5.8 and remained theoretically visible to human eyes for approximately 30 seconds.

September 16, 2008, GRB 080916C discovered by new Fermi Satellite, 12.2 billion light years way, was the intrinsically brightest optical event ever recorded, equivalent to brightness of 9000 supernovae.

April 23, 2009, GRB 090423 discovered by the Swift satellite, the most distant object ever observed in the Universe, about 13.1 billion years ago, when the Universe was only 630 million years old.

If gamma-ray bursts shine equally in all directions, the energy released in gamma rays would be $1000-10,000 \times SN$ or $10-100 \times CO$ core collapse neutrinos.

Comparable to total annihilation of entire star into pure energy!