Monday, February 2, 2015

First exam Friday. First Sky Watch Due.

Review sheet posted Today.

Review session Thursday, 5 – 6 PM, RLM 6.104


Reading:

Chapter 6 Supernovae, Sections § 6.1, 6.2, 6.3 Chapter 1 Introduction, §1.1, 1.2.1, 1.3.1, 1.3.2 Chapter 5 White Dwarfs, § 5.1

Questions? Raise your hand high

Astronomy in the news?

Patriots' goal-line interception



Goal:

To understand the observed nature of supernovae and determine whether they came from white dwarfs or massive stars that undergo core collapse.

Goal:

To understand what we have learned from the study of "live" supernova explosions in other galaxies.

Production of the Elements:

In massive stars (more than about 12 - 15 times the Sun) the core is composed of Helium or heavier elements: Carbon, Oxygen, Magnesium, Silicon, Calcium, finally Iron.

The intermediate-mass elements are produced in the star before the explosion and then expelled into space.

In exploding white dwarfs (arising in stars with mass less than about 8 times the Sun), the core is composed of Carbon and Oxygen, and the explosion creates the intermediate-mass elements, Magnesium, Silicon, Calcium, and also Iron.

(between about 8 and about 12 solar masses, different story, maybe collapsing white dwarfs)

Stellar Physics:

There are many more low mass stars born than high mass stars.

High mass stars have more fuel to burn, but they burn much hotter and brighter. As a result they live a SHORTER time.

A short-lived star must be massive.

A long-lived star must be of relatively low mass.

Galaxy Physics:

Star are born in the spiral arms of spiral galaxies.

Elliptical galaxies have not formed any new stars in billions of years.

Why do the elements carbon, oxygen, magnesium, and silicon frequently appear in the matter ejected from supernovae?




They are built up from the element helium

They are built up from the element calcium

They are built up from the element iron

If a supernova explodes in the spiral arm of a spiral galaxy




It was probably an exploding massive star

It probably ejected silicon

It probably ejected helium

If a supernova explodes far away from any spiral arm of a spiral galaxy



It was probably an exploding massive star

It probably left a neutron star

It probably exploded in an elliptical galaxy

Categories of Supernovae

1st category discovered



Type Ia – near peak light, no detectable Hydrogen or Helium in the spectrum, rather "intermediate mass elements" such as oxygen, magnesium, silicon, sulfur, calcium. Iron appears later as the light fades.

Type Ia occur in all galaxy types:


In spiral galaxies they tend to avoid the spiral arms, they have had time to drift away from the birth site \rightarrow the star that explodes is old

In elliptical galaxies where star formation is thought to have ceased long ago \rightarrow the star that explodes is old, billions of years

⇒the progenitor that explodes must be long-lived, not very massive, suggesting a white dwarf. Sun is long-lived, but won't explode

Type Ia - no hydrogen or helium, intermediate mass elements early, iron later

Light Curve - brightness vs. time consistent with an exploding C/O white dwarf expect total disruption, no neutron star




Type Ia occur in elliptical galaxies, tend to avoid spiral arms in spiral galaxies - old when explode, all evidence points to an exploding white dwarf.


SN 1006, almost definitely Type Ia

Tycho, SN 1572 definitely Type Ia

Recent discovery: spectrum from peak light reflected from surrounding dust, arriving only "now."



Kepler, 1604, probably Type Ia (no sign of neutron star, same ejected composition as SN 1006, Tycho), but some ambiguities.



G1.9+0.3 probably a Type Ia.



Type Ia

no Hydrogen or Helium intermediate mass elements (oxygen, magnesium, silicon, sulfur, calcium) early on, near maximum, iron later avoid spiral arms, occur in elliptical galaxies peaked light curve no neutron star

all consistent with thermonuclear explosion in white dwarf that has waited for a long time (hundreds of millons to billions of years) to explode, total disruption Type II Supernovae - "other" type discovered early in the study of supernovae, show Hydrogen in the spectrum early, Oxygen, Magnesium, Calcium, later


Most occur in spiral galaxies, in the spiral arms, they have no time to drift from the birth site

never in elliptical galaxies (no young stars)

Stars with more mass have more fuel, but they burn it at a prodigous rate, live a shorter time!

→ The progenitor stars are young, short-lived (millions to tens of millions of years) massive stars

We expect such stars to evolve to form iron cores and collapse to a neutron star or black hole (physics to come)



SN 1999em


"Plateau" light curves of Type II are consistent with explosion in a Red Giant

Betelgeuse is a massive red giant, 15 solar masses: we expect it to become a Type II supernova. *Maybe tonight!* Rigel in Orion probably burning He to C/O, explode later.

SN 386, 1181 records are sparse, might have been Type II


Crab was "peculiar" (high helium abundance, slow explosion), but probably a Type II. Expelled outer hydrogen envelope has been difficult to detect directly, but is inferred.

Cas A was probably something else with a very thin layer of Hydrogen (next topic),




Type II are common in other galaxies, more frequent than Type Ia.

Crab nebula







A supernova explodes in an elliptical galaxy. Near peak light what element do you expect to see in the spectrum?









