

Annual Report for Period:09/2007 - 08/2008**Submitted on:** 08/03/2008**Principal Investigator:** Jogee, Shardha .**Award ID:** 0607748**Organization:** U of Texas Austin**Submitted By:**

Jogee, Shardha - Principal Investigator

Title:

Bars and their Impact on Galaxy Evolution over the Last Eight Billion Years

Project Participants**Senior Personnel****Name:** Jogee, Shardha**Worked for more than 160 Hours:** Yes**Contribution to Project:**

9/1/06-8/31/08: Dr. Jogee is the PI of the NSF grant that funded the science project and education and public Outreach (EPO) program. She is responsible for the overall direction and management of the project. She is involved in the scientific analyses and in advising graduate students Irina Marinova, Tim Weinzirl, Amanda Heiderman, postdoctoral fellow Fabio Barazza, and undergraduate Dean Scholars Kyle Penner and Sarah Miller. On the EPO front, Dr Jogee has been responsible for the scientific input and development of the Galaxy & Cosmic Explorer Tool (GCET), working closely with graduate students Achal Augustine and Aaron Smith.

Name: Hemenway, Mary Kay**Worked for more than 160 Hours:** No**Contribution to Project:**

9/1/06-8/31/08: Dr. Hemenway leads the education team and guides the development of the education activities for K-12 students. Dr. Hemenway is supported by this grant and a NASA grant.

Name: Lester, Daniel**Worked for more than 160 Hours:** No**Contribution to Project:**

9/1/06-8/31/07: Dr Lester helped in the preparation of the K-12 activities. In collaboration with the PI , he prepared a proposal to UT for developing an instructional tool for activities on galaxy evolution. Dr. Lester's time is funded by McDonald Observatory.

Name: Preston, Sandra**Worked for more than 160 Hours:** No**Contribution to Project:**

9/1/06-8/31/08: Ms. Preston supervises the Education and Outreach staff, prepares reports and sets up the team meetings. She is supported by the McDonald Observatory

Post-doc**Name:** Barazza, Fabio**Worked for more than 160 Hours:** No**Contribution to Project:**

9/1/06-03/31/07: Dr Barazza is a postdoctoral fellow working on disk galaxies and bars in SDSS and GEMS. He is supported by a NASA grant.

Graduate Student**Name:** Marinova, Irina**Worked for more than 160 Hours:** Yes**Contribution to Project:**

9/1/06-8/31/08: Ms Marinova is a graduate student funded by the NSF project. The proposed program will constitute a major part of her masters and Ph.D.D. thesis.

Name: Worhatch, Randi

Worked for more than 160 Hours: No

Contribution to Project:

6/1/07-8/31/07: Ms. Worhatch, an astronomy graduate student, worked on the formative evaluation to determine what activities would be of most interest to students. She also assisted in the creation of an activity and helped to field test it. Ms. Worhatch was supported by this grant and a NASA grant.

Name: Weinzirl, Tim

Worked for more than 160 Hours: Yes

Contribution to Project:

12/1/07-8/31/08: Worked on bulge-bar-disk decomposition to constrain the origin of bulges in hierarchical models of galaxy evolution. Supported by this grant and several other NASA and startup grants

Name: Heiderman, Amanda

Worked for more than 160 Hours: Yes

Contribution to Project:

06/01/08-8/31/08: Helped to classify bars in the STAGES survey and characterize galaxy interactions. Supported by this grant and several other NASA and startup grants.

Undergraduate Student

Technician, Programmer

Name: Fricke, Kyle

Worked for more than 160 Hours: No

Contribution to Project:

9/1/06-8/31/08: Mr. Fricke prepares the educational activities, field tests them with teachers, and has assisted with the creation of the GCET tool. Mr. Fricke is supported by this grant and a NASA grant.

Other Participant

Research Experience for Undergraduates

Organizational Partners

NASA Virtual Design Center

Other Collaborators or Contacts

I) RESEARCH PROGRAM

Collaborators who are not funded by the NSF grant include Dr Fabio Barazza, Dr Isaac Shlosman, Dr Hans-Walter Rix, Dr Chris Conselice, Dr Chien Peng, Dr Andreas Burkert, Dr Sadegh Khochfar.

II) EPO PROGRAM

Collaborators who are not funded by the NSF grant include the Faculty And Student Teams for Technology (FAST Tex) at the University of Texas Division of Instructional Innovation and Assessment (DIIA), as well as Texas high school teachers Chris Cotter, Christy Budynkiewicz, and Karen Green, who have helped test ideas and the activities in their classrooms. This grant has provided stipends to Cotter and Budynkiewicz.

Activities and Findings

Research and Education Activities:

1. Dr Jogee and her research group at UT Austin have led papers in the areas below.

A) Bars as a function of environment

B) The origin of bulges and the problem of bulgeless galaxies

C) History of Galaxy Interactions and their Impact on SF over the Last 7 Gyr

D) Properties and Impact of Interacting Galaxies in the A901/02 Supercluster from STAGES

2. Furthermore, in collaboration with other co-team members from the GEMS, Coma, STAGES collaboration, Dr. Jogee has worked on and co-authored papers that address the evolution of galaxies as a function of redshift and environment

- 'GEMS Survey Data and Catalog'
- 'An Explanation for the Observed Weak Size Evolution of Disk Galaxies'
- 'The HST/ACS Coma Cluster Survey: I - Survey Objectives and Design'
- 'The dark matter environment of the Abell 901/902 supercluster: a weak lensing analysis of the HST STAGES survey'

II) EPO PROGRAM

A Galaxies and Cosmos Explorer Tool (GCET; <http://www.as.utexas.edu/gcet/>) has been created using the recent large galaxy surveys conducted with NASA's Hubble Space Telescope Advanced Camera for Surveys (ACS). The GCET is an online web-based tool that allows the general public and students to actively participate in quantitative analyses of HST images from the Galaxy Evolution from Morphology and SEDs (GEMS) survey, one of the widest-area galaxy surveys conducted in two filters with ACS to date. The tool allows users to surf the vast cosmos and access ACS images of over 8,000 galaxies over the last eight billion years. For galaxies of interest, users can measure the size, determine the lookback time for concordance cosmology, perform morphological classification on images at two rest-frame wavelengths, and gauge the different stellar populations present. Users can record their measurements, as well as reference information, such as coordinates and redshift, of each galaxy into Excel spreadsheets for further analysis. The celestial coordinates can be used to extract further multiwavelength data from existing archives and upcoming virtual observatories. For undergraduate classes, more advanced IDL or C-based analyses that employ the full samples, can be combined with the visualization capabilities of GCET in order to explore the nature of interesting objects, such as the most massive galaxies, starbursting systems, as well as interacting and merging galaxies. GCET provides a powerful tool for discovery learning in undergraduate introductory science classes as well as high schools.

The testing phases of the tool took place from July to Nov 2007 and a pilot version was released for testing in classrooms in Feb 2008. Final improvements based on feedback were made in summer 2008 for the final version to be released in Sep 2008.

The third activity ('Lives of Stars') was also created by our team. It was pilot-tested with a group of high-school teachers and in a high-school classroom with students. The first three activities were presented to teachers in a 3-hour short course at the Conference for the Advancement of Science Teaching in November, 2007.

The last two activities are in the early stages of development. A six-page student guide for using GCET has been created and field-tested with high-school teachers with positive results. The guide is the fundamental piece in both activities and is currently being further developed and improved upon.

We are working with folks from Google to make GCET available at Google Sky

Findings:

I) RESEARCH PROGRAM

I provide below, under (A) to (D) a summary of the results of my research group, which included over 2007-2008 three beginning graduate

students (A. Heiderman, T. Weinzirl, I. Marinova), two undergraduate Dean scholars (S. Miller and K. Penner), and postdoctoral fellow F. Barazza.

A) Bars as a function of environment

- Barazza, F.-D., Jablonca, P., Desai, V., Jogee, S. & the EDisCS collaboration 2008
- Marinova, I., Jogee, S., & the STAGES collaboration 2008a
- Marinova, I., Jogee, S., & the STAGES collaboration 2008b
- Weinzirl, T., Jogee, S. & the Coma collaboration, in prep

While bars in the field have been widely studied, comparatively little is known about the frequency, properties, and impact of bars in rich clusters. Not only do clusters provide an interesting laboratory to test bar formation models, but bars can also be used to test the mode of cluster growth. Using the bar analysis package and quantitative approach established in Jogee et al (2004), we are currently exploring bars in clusters through two studies: a study of 800 bright galaxies in the STAGES A901/902 supercluster survey at a redshift $z \sim 0.17$. (Marinova, Jogee, & the STAGES collaboration 2008b), as well as a study of 500 galaxies in the ACS treasury survey of the rich Coma cluster at $z \sim 0.025$ (Weinzirl, Jogee, & the Coma collaboration, in prep.). Dr Jogee is also collaborating on a related study of 2256 disk galaxies in the EDisCS survey of clusters at $z \sim 0.4$ to 1.0 (Barazza, Jablonka, Desai, Jogee & the EDisCS collaboration 2008b).

Our early results from the STAGES survey suggest that the optical bar fraction in the rich A901 and A902 clusters is similar to that of the field, and shows no significant trend with any local environment tracer, such as the projected mass density κ , Σ_{10} , ICM density from X-ray emission, and the projected distance to the nearest cluster center (Marinova, Jogee, & the STAGES collaboration 2008). Similarly, no significant difference is found between the optical bar fraction of field and clusters over $z \sim 0.4$ to 1.0 in the EDisCS survey (Barazza et al 2008b). The latter study also finds no evidence for any strong decline in the optical bar fraction with redshift. Taken together, our results increasingly suggest that the processes controlling the frequency and properties of bars are not a strong function of environment.

B) The origin of bulges and the problem of bulgeless galaxies

- Weinzirl, T. Jogee, S., & Barazza, F. 2008
- Weinzirl, T., Jogee, S. Khochfar, S., Burkert, A., & Kormendy, J. 2008 (WJKBK08)

In Lambda CDM models of galaxy evolution, there are in principle three main mechanisms to build bulges of spiral galaxies: major mergers, minor mergers, and secular processes (see WJKBK08 for details). The major merger of two spiral galaxies destroys the disk component and leaves behind a classical bulge, around which a stellar disk forms when hot gas in the halo subsequently cools, settles into a disk, and forms stars. Minor mergers can also grow bulges in several ways. A tidally induced bar and/or direct tidal torques from the companion can drive gas into the inner kpc (e.g., Quinn et al. 1993; Hernquist & Mihos 1995; Jogee 2006 and references therein), where subsequent SF forms a compact high v/σ stellar component, or disk pseudobulge. In addition, the stellar core of the satellite can sink to the central region via dynamical friction. Finally, bulges can also have a secular origin: here, a stellar bar or globally oval structure in a **non-interacting** galaxy drives gas inflow into the inner kpc, where subsequent SF forms a disk pseudobulge (e.g., Kormendy 1993; Jogee 1999; Kormendy & Kennicutt 2004; Jogee, Scoville, & Kenney 2005).

These different mechanisms to form bulges have been postulated for a long time. However, what is still missing is a quantitative assessment of the relative importance of different bulge formation pathways in high and low mass spirals. For instance, although bulges are an integral part of massive present-day spiral galaxies, we still cannot answer the following basic question: do most bulges in massive spirals form via major mergers, minor mergers, or secular processes?

In WJKBK08, we attempt one of the first quantitative comparisons of the properties of bulges in a fairly complete sample of high mass ($M^* \geq 10^{10}$ Mo) spirals to predictions from Lambda CDM-based simulations of galaxy evolution. We derive the bulge-to-total mass ratio (B/T) and bulge Sérsic index n by performing 2D bulge-disk-bar decomposition on H-band images of 146 bright, high mass, moderately inclined spirals.

1) Interestingly, we find that as many as ~56% of high mass spirals have low $n \leq 2$ bulges: such bulges exist in barred and unbarred galaxies across all Hubble types. Furthermore a striking ~66% of high mass spirals have $B/T \leq 0.2$

2) We compare the observed distribution of bulge B/T in high mass spirals to predictions from Λ -CDM-based semi-analytical models. In the models, a bulge with $B/T \leq 0.2$ can exist in a galaxy with a past major merger, only if the last major merger occurred at $z > 2$ (lookback > 10 Gyr). The predicted fraction of high mass spirals with a past major merger and a bulge with a present-day $B/T \leq 0.2$ is a factor of over fifteen smaller than the observed fraction (~66%) of high mass spirals with $B/T \leq 0.2$. The comparisons rule out major mergers as the main formation pathway for bulges in high mass spirals. Contrary to common perception, bulges built via major mergers seriously fail to account for the bulges present in ~66% of high mass spirals. !

3) In the models, the majority of low $B/T \leq 0.2$ bulges exist in systems that have experienced only minor mergers, and no major mergers. These bulges can be built via minor mergers and secular processes. So far, we explored one realization of the model focusing on bulges built via satellite stars in minor mergers and find good agreement with the observations. Future models will explore more realistic minor merger scenarios and secular processes in paper~II.

C) History of Galaxy Interactions and their Impact on SF over the Last 7 Gyr

- Jogee, S, et al. & the GEMS collaboration, 2008a
- Jogee, S, et al. & the GEMS collaboration, 2008b
- Miller, S., Jogee S, & the GEMS collaboration 2008

The merger history of galaxies impacts the mass assembly, star formation history, AGN activity, and structural evolution of galaxies. The merger rate/fraction at $z > 1$ remains highly uncertain, owing to relatively modest volumes and bandpass shifting effects, but with a general trend towards higher merger fractions at higher redshifts. Even the merger rate at $z < 1$ has proved hard to robustly measure for a variety of reasons, ranging from small samples in early studies, to different methods on large samples in later studies.

In Jogee et al. (2008a,b), we have performed a complementary and comprehensive observational estimate of the frequency of interacting galaxies over $z \sim 0.24$ – 0.80 (lookback times of 3–7 Gyr), and the impact of interactions on the SF of galaxies over this interval. Our study is based on HST ACS, COMBO-17, and Spitzer 24 μ m data from the GEMS survey. We use a large sample of ~ 3600 ($M > 1e9$ Mo) galaxies and ~ 790 high mass ($M > 2.5e10$ Mo) galaxies for robust number statistics. Two independent methods are used to identify strongly interacting galaxies: a tailored visual classification system complemented with spectrophotometric redshifts and stellar masses, as well as the CAS merger criterion ($A > 0.35$ and $A > S$; Conselice 2003). This allows one of the most extensive comparisons to date between CAS-based and visual classification results. While many earlier studies focused only on major mergers, we try to constrain the frequency of minor mergers as well, since they dominate the merger rates in Λ -CDM models. Some of our results are outlined below.

1) Among ~ 790 high mass galaxies, the fraction of visually-classified interacting systems over lookback times of 3–7 Gyr ranges from $9\% \pm 5\%$ at $z \sim 0.24$ – 0.34 , to $8\% \pm 2\%$ at $z \sim 0.60$ – 0.80 , as averaged over every Gyr bin. These systems appear to be in merging or post-merger phases, and are candidates for a recent merger of mass ratio $M1/M2 > 1/10$. The lower limit on the major ($M1/M2 > 1/4$) merger fraction ranges from 1.1% to 3.5% over $z \sim 0.24$ – 0.80 . The corresponding lower limit on the minor ($1/10 \leq M1/M2 < 1/4$) merger fraction ranges from 3.6% to 7.5%. This is the first, albeit approximate, empirical estimate of the frequency of minor mergers over the last 7 Gyr.

For an assumed value of ~ 0.5 Gyr for the visibility timescale, it follows that each massive ($M > 2.5e10$ Mo) galaxy has undergone ~ 0.7 mergers of mass ratio $> 1/10$ over the redshift interval $z \sim 0.24$ – 0.80 . Of these, we estimate that $1/4$ are major mergers, $2/3$ are minor mergers, and the rest are ambiguous cases of major or minor mergers. The corresponding merger rate R is a few times $1e-4$ galaxies Gyr $^{-1}$ Mpc $^{-3}$. Among ~ 2840 blue cloud galaxies of mass $M > 1e9$ Mo, similar results hold.

2) We compare our empirical merger rate R for high mass galaxies to predictions from different Λ CDM-based simulations of galaxy evolution, including the halo occupation distribution (HOD) models of Hopkins et al (2007); semi-analytic models (SAMs) of Somerville et al. (2008), Bower et al. (2006), and Khochfar & Silk (2006); and smoothed particle hydrodynamics (SPH) cosmological simulations from Maller et al. (2006). To our knowledge, such extensive comparisons have not been attempted to date, and are long overdue. We find qualitative agreement between the observations and models, with the (major+minor) merger rate from different models bracketing the observed rate, and

showing a factor of five dispersion.} One can now anticipate that in the near future, improvements in both the observational estimates and model predictions will start to rule out certain merger scenarios and refine our understanding of the merger history of galaxies.

3) The idea that galaxy interactions generally enhance the SFR of galaxies is well established from observations (e.g., Joseph & Wright 1985; Kennicutt et al. 1987), and simulations (e.g., Hernquist 1989; Mihos & Hernquist 1994, 1996; Springel, Di Matteo & Hernquist 2005b). However, simulations cannot uniquely predict the factor by which interaction enhance the SF activity of galaxies over the last 7 Gyr, since both the SFR and properties of the remnants in simulations are highly sensitive to the stellar feedback model, the bulge-to-disk (B/D) ratio, the gas mass fractions, and orbital geometry (e.g., Cox et al 2006; di Matteo et al. 2007). Thus, empirical constraints are needed. Among ~ 3600 $M \geq 1e9$ Mo galaxies, we find that the average SFR of visibly interacting galaxies is only modestly enhanced compared to non-interacting galaxies over $z \sim 0.24 - 0.80$. This result is found for SFRs based on UV, UV+IR, and UV+stacked-IR data. This modest enhancement is consistent with the results of di Matteo et al. (2007) based on numerical simulations of several hundred galaxy collisions.

4) The SF properties of interacting and non-interacting galaxies since $z < 1$ are of great astrophysical interest, given that the cosmic SFR density is claimed to decline by a factor of 4 to 10 since $z \sim 1$ (e.g., Lilly et al. 1996; Ellis et al 1996; Hopkins 2004; Pérez-González et al. 2005; Le Floc'h et al. 2005). We therefore set quantitative limits on the contribution of obviously interacting systems to the UV-based and UV+IR-based SFR density over $z \sim 0.24 - 0.80$. Among ~ 3600 $M \geq 1e9$ Mo galaxies, we find that visibly interacting systems only account for a small fraction ($< 30\%$) of the cosmic SFR density over lookback times of $\sim 3 - 7$ Gyr ($z \sim 0.24 - 0.80$). In effect, our result suggests that the behavior of the cosmic SFR density over the last 7 Gyr is predominantly shaped by non-interacting galaxies, rather than strongly interacting galaxies. This suggests that the observed decline in the cosmic SFR density since $z \sim 0.80$ is largely the result of a shutdown in the SF of non-interacting galaxies.

D) Properties and Impact of Interacting Galaxies in the A901/02 Supercluster from STAGES

- Heiderman, A., Jogee, S., & the STAGES collaboration 2008a

- Heiderman, A., Jogee, S., & the STAGES collaboration 2008b

In project (C), we looked at the history and impact of galaxy interaction in the *field* over the last 7 Gyr. In project (D), we extend this work from field to three rich galaxy clusters within the Abell 901/902 supercluster at $z \sim 0.165$. We present a study of interacting galaxies and the influence of environment in the A901/02 supercluster, based on 774 bright, intermediate mass ($M_V < -18$; $M > 1e9$ Mo) galaxies. We use HST ACS F606W data from the STAGES survey, COMBO-17, Spitzer 24 micron, and XMM-Newton X-ray data. We use visual classification, as well as quantitative CAS parameters to estimate the fraction f_{int} of strongly interacting galaxies, which are likely candidates for interactions of stellar mass ratio $M_1/M_2 > 1/10$. Our findings are as follows

(1) We find, based on visual classification, that the fraction (f_{int}) of strongly interacting galaxies is 0.049 ± 0.013 , with at least 0.015 ± 0.005 being major interactions ($1/4 < M_1/M_2 \leq 1$), at least 0.012 ± 0.005 being minor interactions ($1/10 < M_1/M_2 \leq 1/4$), and 0.022 ± 0.007 being ambiguous cases of major or minor interactions. We compare f_{int} to groups and clusters at other redshifts.

(2) The strongly interacting galaxies lie outside the cluster core in the region between the core and virial radius, at clustocentric radii $0.25 \text{ Mpc} < R \leq 1.2 \text{ Mpc}$. We suggest that this is due to the velocity dispersion of cluster galaxies falling outside the core, or/and due to groups being accreted by the A901/902 clusters. Our number and distribution of strong interactions shows fair agreement with those predicted by N-body simulations of accreting groups in the A901/902 clusters.

(3) The fraction f_{blue} of bright galaxies that lie on the blue rises from 0.09 ± 0.03 , to 0.41 ± 0.05 as we move from the cluster core to the virial radius. For strongly interacting galaxies, f_{blue} is 0.61 ± 0.14 (23/38) versus 0.33 ± 0.07 (257/774) for non-interacting galaxies, implying that strongly interacting galaxies are preferentially blue.

(4) The average SFR, based on UV and IR data, is enhanced by a factor of ~ 2 to 4 in strongly interacting galaxies compared to non-interacting galaxies. However, strongly interacting galaxies only contributes 20% of the total SFR density in the A901/902 clusters. The remaining SFR density comes from galaxies whose SFR is significantly depressed at a given stellar mass compared to field galaxies of similar redshifts. The SFR is depressed in the core ($R \leq 0.25 \text{ Mpc}$), but also out to the virial radius ($R \leq 1.2 \text{ Mpc}$). This implies that the processes responsible for the depression must be effective over a large range of radii.

II) EPO PROGRAM

In the process of planning the educational activities, we were introduced to the NASA Virtual Design Center (VDC), a service of the Mid-Atlantic Region Space Science Broker, at the Astronomical Society of the Pacific meeting in September 16-18, 2006. Using the VDC design template, our team began by conducting a survey of high school students to determine their current level of astronomical knowledge and to identify the areas of astronomy that interest them most.

Shortly after the survey was conducted, on October 9, we contacted Dr. Laurie Ruberg at the VDC to formally request facilitation. The education team met with Dr. Ruberg and her staff by telecon on October 13, December 13, and May 8, and by videoconference on November 17. In addition, Dr. Hemenway met with Dr. Ruberg for two hours in Seattle on January 6, 2007 when both were attending the AAS/AAPT national meeting.

The VDC has provided written assessment of our product. Their suggestions have been of great value in developing our product, including both implementation and assessment. It is our intention to continue using their design process as we continue with the other activities that complete the package for this project.

During the original planning meeting, we identified an opportunity to submit a proposal to the FAST Tex program at the University of Texas Division of Instructional Innovation and Assessment (DIIA). As outlined above, the PI and Dr. Lester submitted a successful proposal, which was awarded 300 student hours to create the Galaxies and Cosmos Explorer Tool (GCET) for undergraduates and K-12 students. The PI has worked with two computer science and DIIA graduate students (Achal Augustine and Aaron Smith) to develop GCET. The testing of GCET took place from July to Nov 2007 and a pilot version released for testing in classrooms in Feb 2008. Final improvements based on feedback are being made in summer 2008 for the final version to be released in Sep 2008. The tool is described in detail under the section 'Publications and Products'.

Training and Development:

I) RESEARCH PROGRAM

Graduate students I. Marinova, T. Weinzirl, A. Heiderman, and postdoctoral fellow Barazza have developed proficiency in programming with the Interactive Data Language (IDL); statistical analyzes; isophotal analyzes of images with the IRAF package; 2D bulge-disk-bar structural decomposition with Python scripts and the GALFIT software, and a deeper understanding of the orbital structure of rotating barred potentials. They have also developed better communication skills through multiple oral presentations of their research in seminars and at conferences.

II) EPO PROGRAM

While developing GCET with the PI, graduate students Achal Augustine (in computer science) and Aaron Smith (in instructional technologies) learn about research in astronomy based on large galaxy surveys conducted with HST, and developed new skills in programming with Flash, Ajax, and Java for instructional innovation.

Outreach Activities:

A short course entitled Evolving Galaxies/Evolving Science was presented to high school teachers at the Conference for the Advancement of Science Teaching in Austin, Texas on November 17, 2007. Another short course has been proposed for the November 2008 meeting in Fort Worth, Texas.

Journal Publications

Marinova, I. & Jogee, S, "Characterizing Bars at $z \sim 0$ in the optical and NIR: Implications for the Evolution of Barred Disks with Redshift", ApJ, p. 1176, vol. 659, (2007). Published,

Barazza, F., Jogee, S., & Marinova, I, "Bars in Disk-Dominated and Bulge-Dominated Galaxies at $z \sim 0$: New Insights from 3600 SDSS Galaxies", *ApJ*, p. 1194, vol. 675, (2008). Published,

Barazza, F.~D.; Jablonca, P.; Desai, V.; Jogee, S; and the ESO Distant Clusters Survey Collaboration, "Frequency and properties of bars in cluster and field galaxies at intermediate redshifts from EDisCS", *Astrophysical Journal*, p. , vol. , (2008). Submitted,

Caldwell, JAR; McIntosh, DH; Rix, HW; Barden, M; Beckwith, SVW; Bell, EF; Borch, A; Heymans, C; Haussler, B; Jahnke, K; Jogee, S; Meisenheimer, K; Peng, CY; Sanchez, SF; Somerville, RS; Wisotzki, L; Wolf, C, "Gems survey data and catalog", *ASTROPHYSICAL JOURNAL SUPPLEMENT SERIES*, p. 136, vol. 174, (2008). Published,

Carter, D., et al., "The HST/ACS Coma Cluster Survey: I - Survey Objectives and Design", *Astrophysical Journal*, p. , vol. , (2008). Published,

Jogee, S. et al., "History of Galaxy Interactions and Their Impact on Star Formation over the Last 7 Gyr from GEMS", *Astrophysical Journal*, p. , vol. , (2008). Submitted,

Weinzirl, T., Jogee, S.; Khochfar, S.; Burkert, A.; Kormendy, J. 2008, "Bulge σ and B/T in High Mass Galaxies: Constraints on the Origin of Bulges in Hierarchical Models", *Astrophysical Journal*, p. , vol. , (2008). Submitted,

Gallazi, A. et al., "Obscured Star Formation in Intermediate Density Environment", *Astrophysical Journal*, p. , vol. , (2008). Submitted,

Somerville, R. S. et al., "An Explanation for the Observed Weak Size Evolution of Disk Galaxies", *Astrophysical Journal*, p. , vol. , (2008). Published,

Heymans, C., et al., "The dark matter environment of the Abell 901/902 supercluster: a weak lensing analysis of the HST STAGES survey", *MNRAS*, p. , vol. , (2008). Published,

Heiderman, A.; Jogee, S; the STAGES collaboration, "Properties and Impact of Interacting Galaxies in the Abell 901/902 Supercluster from STAGES", *Astrophysical Journal*, p. , vol. , (2008). In preparation (submission by Aug 10/2008),

Marinova, I.; Jogee, S; the STAGES collaboration, "The Properties of Barred Disks in a Supercluster Environment: Constraints from Abell 901/2 with STAGES", *Astrophysical Journal*, p. , vol. , (2008). In preparation (submission by Aug 10/2008),

Books or Other One-time Publications

Barazza, F.~D., Jogee, S., & Marinova, I, "Constraints on Bars in the Local Universe from 5000 SDSS Galaxies", (2007). Book, Published

Editor(s): F. Combes and J. Palous

Collection: Galaxy Evolution Across the Hubble Time, Proceedings of the International Astronomical Union 2, IAU Symposium #235

Bibliography: Barazza, F.~D., Jogee, S., & Marinova, I. 2007, IAU Symposium, 235, 76

Penner, K. Jogee, S., Miller, S., and GEMS collaboration, "Constraining the interaction history of galaxies over 8 Gyr", (2007). Book, Published

Collection: BAAS

Bibliography: Penner, K., Jogee, S., Miller, S., & GEMS collaboration, 2007, BAAS

Hemenway, Mary Kay; Jogee, S; Augustine, A; Smith, A; and Lester, D, "The Galaxy and Cosmos Explorer Tool", (2007). Book, Submitted
Collection: 2007 COSMOS in the Classroom Meeting on Aug 2-5, 2007

Bibliography: Hemenway, Mary Kay; Jogee, S; Augustine, A; Smith, A; and Lester, D.; The Galaxy and Cosmos Explorer Tool (abstract submitted to the 2007 COSMOS in the Classroom Meeting on Aug 2-5

Hemenway, Mary Kay; Jogee, S.; Fricke, K.; Worhatch, R.; Ruberg, L. F, "Developing the 'Multiwavelength Astronomy: Galaxies in a Different Light' Activity

", (2007). Book, Submitted

Collection: 2007 ASP Annual Conference on Sep 5-7, 2007

Bibliography: Hemenway, Mary Kay; Jogee, S.; Fricke, K.; Worhatch, R.; Ruberg, L. F.; Developing the ?Multiwavelength Astronomy: Galaxies in a Different Light? Activity (abstract submitted to t

Smith, A., Rhodes, S., and Jogee, S., "The Galaxy and Cosmos Explorer Tool", (2007). Book, Submitted

Collection: Instructional Technology showcase <http://www.utexas.edu/academic/diia/itshowcase/>)

Bibliography: Smith, A., Rhodes, S., and Jogee, S., 'The Galaxy and Cosmos Explorer Tool' (abstract submitted for the 2007 Instructional Technology showcase)

Jogee, S. et al. 2008, "Frequency and Impact of Galaxy Mergers and Interactions over the last 7 Gyr", (2008). Book, Accepted

Editor(s): J.~G.~Funes, S.~J., & E.~M.~Corsini (San Francisco: ASP)

Collection: Formation and Evolution of Galaxy Disks

Bibliography: Jogee, S. et al. 2008, in Formation and Evolution of Galaxy Disks, ed.~J.~G.~Funes, S.~J., & E.~M.~Corsini (San Francisco: ASP), in press (arXiv:0802.3901)

Jogee, S; Hemenway, M K; Miller, S; Smith, A; Augustine, A; Fricke, K; Worhatch, R; Preston, S; and Lester, D., "The Galaxies and Cosmos Explorer Tool: Charting Galaxies Over Cosmic Times in the Classroom", (2008). 211th AAS meeting BAAS, Accepted

Collection: 211th AAS meeting BAAS

Bibliography: BAAS, 211, 06.12

Jogee, S., "Evolution of Disk Galaxies: New Insights and Future Challenges (Invited Plenary Lecture)", (). 211th AAS meeting BAAS, Accepted

Editor(s): 211th AAS meeting BAAS

Bibliography: BAAS 211, 88.01

Jogee, S., et al., "Star Formation in Interacting and Normal Galaxies over the last 7 Gigayear.", (2008). 211th AAS meeting BAAS, Accepted

Collection: 211th AAS meeting BAAS

Bibliography: BAAS, 211, 126.06

Marinova, I.; Jogee, S; the STAGES collaboration, " Characterizing Barred Galaxies in Abell 901/902", (2008). Book, Accepted

Editor(s): A. Frebel, J. Maund, J. Shen, M. Siegel

Collection: ASP conference proceedings of the Frank N. Bash Symposium 2007: New Horizons in Astronomy

Bibliography: Marinova, I., Jogee, S., & the STAGES collaboration, 2008, ASP conference proceedings of the "Frank N. Bash Symposium 2007: New Horizons in Astronomy", Editors A. Frebel, J. Mau

Miller, S.; Jogee, S.; and the GEMS collaboration

, "Exploring the Impact of Galaxy Interactions over Seven Billion Years with CAS", (2008). Book, Accepted

Editor(s): A. Frebel, J. Maund, J. Shen, M. Siegel

Collection: ASP conference proceedings of the Frank N. Bash Symposium 2007: New Horizons in Astronomy

Bibliography: Miller, S., Jogee, S., and the GEMS collaboration 2008, ASP conference proceedings of the "Frank N. Bash Symposium 2007: New Horizons in Astronomy", Editors A. Frebel, J. Maund, J

Heiderman, A.; Jogee, S; the STAGES collaboration, "Galaxy Evolution in the A901/02 Supercluster: Constraints from Galaxy-Galaxy and Galaxy-ICM Interactions.", (2008). Book, Accepted

Editor(s): A. Frebel, J. Maund, J. Shen, M. Siegel

Collection: ASP conference proceedings of the Frank N. Bash Symposium 2007: New Horizons in Astronomy

Bibliography: Heiderman, A., Jogee, S., & the STAGES collaboration, 2008, ASP conference proceedings of the "Frank N. Bash Symposium 2007: New Horizons in Astronomy", Editors A. Frebel, J. Mau

Weinzirl, T., Jogee, S; Barazza, F., "Constraining Galaxy Evolution With Bulge+Disk+Bar Decomposition", (). 211th AAS meeting BAAS, Accepted

Collection: 211th AAS meeting BAAS

Bibliography: BAAS, 211, 97.09

Weinzirl, T., Jogee, S; Barazza, F., "Constraints on Disk and Bulge Assembly from Structural Decomposition", (2008). Book, Accepted
 Editor(s): A. Frebel, J. Maund, J. Shen, M. Siegel
 Collection: ASP conference proceedings of the Frank N. Bash Symposium 2007: New Horizons in Astronomy
 Bibliography: Weinzirl, T., Jogee, S, & Barazza, F. 2008, ASP conference proceedings of the "Frank N. Bash Symposium 2007: New Horizons in Astronomy", Editors A. Frebel, J. Maund, J. Shen, M. S

Web/Internet Site

URL(s):

<http://www.as.utexas.edu/gcet/>

Description:

Website 1 featureS the Galaxy and Cosmos Explorer Tool (GCET). The testing phases of the tool took place from July to Nov 2007 and a pilot version released for testing in classrooms in Feb 2008. Final improvements based on feedback are being made in summer 2008 for the final version to be released in Sep 2008.

Other Specific Products

Product Type:

Teaching aids

Product Description:

The PI has been working with two computer science and DIAA graduate students (Achal Augustine and Aaron Smith) to develop the Galaxy and Cosmos Explorer Tool (GCET). We are in the final phases of testing in June and July and we will release the tool in August 2007. GCET is an online web-based tool that allows students to actively engage in the exciting adventure of exploring the evolution of galaxies over a large fraction of the age of the Universe. GCET allows students to perform quantitative analyses of HST images from the Galaxy Evolution from Morphology and SEDs (GEMS) survey, one of the widest-area galaxy surveys conducted with HST to date. The tool allows students to access HST images of over 8,000 galaxies over the last eight billion years, an interval covering two thirds of the age of the Universe. Students can surf the vast cosmos and for galaxies of inte, they can measure the size, determine the lookback time for concordance cosmology, perform morphological classification on images at two rest-frame wavelengths, and gauge the different stellar populations present. Students can record their measurements, as well as reference information (such as coordinates and redshift) of each galaxy into spreadsheets for further analysis. The celestial coordinates can be used to extract further multiwavelength data from existing archives and upcoming virtual observatories. GCET provides a powerful tool for discovery learning in undergraduate science and introductory classes, as well as high schools.

Sharing Information:

The Galaxy and Cosmos Explorer Tool (GCET) is an online web-based tool, whose final version will be released in Sep 2008

Contributions

Contributions within Discipline:

The results described under the section 'Activities and Findings' have advanced our knowledge in the principal disciplinary field of Astronomy, as well as created inquiry-based activities for high school students and undergraduates

Contributions to Other Disciplines:

The creation of the Galaxy and Cosmos Explorer Tool (GCET) has contributed to knowledge of graduate students Achal Augustine (in computer science) and Aaron Smith (in instructional technologies).

Contributions to Human Resource Development:

The activities undertaken under this grant have contributed to the knowledge of the following people:

- (1) Astronomy graduate students I. Marinova, T. Weinzirl, A. Heiderman, and postdoctoral fellow F. Barazza, who worked on the research activities
- (2) Astronomy graduate student, Randi Worhatch, who worked on the EPO activity on multi-wavelength light
- (3) Graduate students Achal Augustine (in computer science) and Aaron Smith (in instructional technologies) who helped to develop 'The

Galaxy and Cosmos Explorer Tool'

Contributions to Resources for Research and Education:

A new tool (GCET) has been developed that will provide undergraduate students and K-12 students with the ability to use cutting-edge data from space and ground-based telescopes to conduct a research project.

Contributions Beyond Science and Engineering:

We are currently having discussions with Google Sky about offering the products produced with this grant at their website.

Special Requirements

Special reporting requirements: None

Change in Objectives or Scope: None

Animal, Human Subjects, Biohazards: None

Categories for which nothing is reported: