



**Astro 358/Spring 2012**



## **Galaxies and the Universe**

Figure for Lecture 2 : Overview of Pre-Requisite Material

### *Pre-requisite for Galaxies Class: Understand stars first!*

- A galaxy is a collection of stars, gas, dust, and dark matter, which are bound by gravity. It contains a few  $10^8$  to  $10^{12}$  stars, which orbit a common center.



HST image of the barred spiral galaxy NGC 1300




Giant elliptical M87



Peculiar/Interacting galaxies  
HST image of The Antennae

### Wien's Law: Surface Temperature and Color of Stars




**Wien's law:** A star or blackbody will emit its maximum flux at a wavelength  $\lambda_{\text{peak}}$  that depends inversely on its surface temperature  $T$


$$\lambda_{\text{peak}} = W / T, \text{ where } W = \text{Wien's constant} = 2.9 \times 10^{-3} \text{ m K}$$

Wien's law implies that hotter stars have smaller peak wavelengths  $\lambda_{\text{peak}}$

### Implication of Wien's Law for Galaxy Images



X-ray/ROSAT



Ultraviolet/ASTRO-1



Visible light



Near infrared/Spitzer

- UV images: mainly trace very hot ( $T > 10^5$  K) massive, short-lived young stars
- Visible images: traced moderately hot ( $T = 5800$  K), moderate mass longer-lived stars like the Sun (lifetime of 10 billion years)
- Near IR : traces cooler ( $T \sim 3000$  K) low mass, long-lived stars AND penetrates through dust revealing these stars
- Mid-IR: traces warm dust ( $T \sim$  few 100 K)

### Dust Obscures Stars and is heated by Massive Hot Stars




Movie  
(NASA/Spitzer)

The optical image of M81 shows intermediate age stars and patches of dusty obscuration.

In the infrared observations of M81 from the Spitzer satellite:

- near-IR light comes from cool ( $T \sim 3000$  K) old stars and it penetrates through dust
- mid-IR light comes warm gas and dust ( $T \sim$  few 100 K) which look clumpy/filamentary

***Hertzsprung-Russell (H-R) diagram:***



Luminosity of a star  
is proportional to  
 $T^4 R^2$

***Mk Classification System For Stars***

*MK = (Spectral Type , Luminosity Class)*  
*OBAFGKM I to V*

*I = Supergiants*

*II = Bright Giants*

*III = Normal Giant*

*IV = Subgiant*

*V = Dwarfs = Main Sequence Stars*

## Original description of spectral types OBAFGKM

**Table 3.1** Principal characteristics of spectral types

Spectral type	Spectral features
O	He II lines visible; lines from highly ionized species, for example, C III, N III, O III, Si IV; H lines relatively weak; strong UV continuum
B	He I lines strong, attain maximum at B2; He II lines absent; H lines stronger; lower-excitation ions, for example, CII, O II, Si III
A	H lines attain maximum strength at A0 and decrease towards later types; Mg II, Si II strong; Ca II weak and increasing in strength
F	H weaker, Ca II stronger; lines of neutral atoms and first ionization states of metals appear prominently
G	Solar-type spectra; Ca II lines extremely strong; neutral metals prominent, ions weaker; G band (CH) strong; H lines weakening
K	Neutral metallic lines dominate; H quite weak; molecular bands (CH, CN) developing; continuum weak in blue
M	Strong molecular bands, particularly TiO; some neutral lines, for example, CA I, quite strong; red continuum
C	Carbon stars; strong bands of carbon compounds C <sub>2</sub> , CN, CO; TiO absent; temperatures in range types K and M
S	Heavy-element stars; bands of ZrO, YO, LaO; neutral atoms strong as in types K and M; overlaps these types in temperature range

From GA

## Relationship between Spectral type and Surface Temperature

**Table 3.7** The effective-temperature and bolometric-correction scales

Spectral type	Luminosity Class					
	V		III		I	
	$T_{\text{eff}}/\text{K}$	$BC_V$	$T_{\text{eff}}/\text{K}$	$BC_V$	$T_{\text{eff}}/\text{K}$	$BC_V$
O3	52 500	-4.75	50 000	-4.58	47 300	-4.41
O5	44 500	-4.40	42 500	-4.05	40 300	-3.87
O7	38 000	-3.68	37 000	-3.58	35 700	-3.48
O9	33 000	-3.33	32 000	-3.13	32 600	-3.18
B0	30 000	-3.16	29 000	-2.88	26 500	-2.49
B2	22 000	-2.35	20 300	-2.02	18 500	-1.58
B3	18 700	-1.94	17 100	-1.60	16 200	-1.26
B5	15 400	-1.46	15 000	-1.30	13 600	-0.95
B7	13 000	-1.02	13 200	-0.97	12 200	-0.78
B8	11 900	-0.80	12 400	-0.82	11 200	-0.66
A0	9 520	-0.30	10 100	-0.42	9 730	-0.41
A5	8 200	-0.15	8 100	-0.14	8 510	-0.13
F0	7 200	-0.09	7 150	-0.11	7 700	-0.01
F5	6 440	-0.14	6 470	-0.14	6 900	-0.03
G0	6 030	-0.18	5 850	-0.20	5 550	-0.15
G2	5 860	-0.20	5 450	-0.27	5 200	-0.21
G5	5 770	-0.21	5 150	-0.34	4 850	-0.33
K0	5 250	-0.31	4 750	-0.50	4 420	-0.50
K5	4 350	-0.72	3 950	-1.02	3 850	-1.01
M0	3 850	-1.28	3 800	-1.25	3 650	-1.29
M5	3 240	-2.73	3 330	-2.48	2 800	-3.47
M8	2 640	-4.1				

From GA

SOURCE: From data published in Schmidt-Kaler (1982)

**Main Sequence Stars**  
(Mass, Luminosity, Absolute Magnitude, Radius)

**Table 3.13** Physical properties of MS stars

Spectral type	$M/M_{\odot}$	$\log(L/L_{\odot})$	$M_{\text{bol}}$	$M_V$	$R/R_{\odot}$	$\bar{\rho}/\bar{\rho}_{\odot}$
O3	120	6.15	-10.7	-6.0	15	0.035
O5	60	5.90	-10.1	-5.7	12	0.035
O8	23	5.23	-8.4	-4.9	8.5	0.037
B0	17.5	4.72	-7.1	-4.0	7.4	0.043
B3	7.6	3.28	-3.5	-1.6	4.8	0.069
B5	5.9	2.92	-2.7	-1.2	3.9	0.099
B8	3.8	2.26	-1.0	-0.2	3.0	0.14
A0	2.9	1.73	0.3	0.6	2.4	0.21
A5	2.0	1.15	1.7	1.9	1.7	0.41
F0	1.6	0.81	2.6	2.7	1.5	0.47
F5	1.3	0.51	3.4	3.5	1.3	0.59
G0	1.05	0.18	4.2	4.4	1.1	0.79
G5	0.92	-0.10	4.9	5.1	0.92	1.18
K0	0.79	-0.38	5.6	5.9	0.85	1.29
K5	0.67	-0.82	6.7	7.4	0.72	1.79
M0	0.51	-1.11	7.4	8.8	0.60	2.36
M5	0.21	-1.96	9.6	12.3	0.27	10.7
M7	0.12	-2.47	10.8	14.3	0.18	20.6
M8	0.06	-2.92	11.9	16.0	0.1	60

From GA

SOURCE: Data published in Schmidt-Kaler (1982)

**Mean V-band magnitudes and colors for main sequence stars**

**Table 1.3** Average magnitudes and colors for main-sequence stars: class V (dwarfs)

	$M_V$	BC	$J5 - V$	$U - B$	$B - V$	$V - R$	$V - I$	$J - K$	$V - K$	$T_{\text{eff}}$
O3	-6	4.5	—	-1.22	-0.32	—	—	—	—	50 000
O5	-5.6	4.0	—	-1.19	-0.32	-0.14	-0.32	-0.25	-0.99	43 000
O8	-4.8	3.3	-4.1	-1.14	-0.32	-0.14	-0.32	-0.24	-0.96	35 000
B0	-4.0	2.9	-4.0	-1.07	-0.30	-0.13	-0.30	-0.23	-0.91	29 800
B3	-1.4	1.6	-2.9	-0.75	-0.18	-0.08	-0.20	-0.15	-0.54	18 750
B6	-1.0	1.2	-2.3	-0.50	-0.14	-0.06	-0.13	-0.09	-0.39	14 000
B8	-0.25	0.8	-1.7	-0.30	-0.11	-0.04	-0.09	-0.06	-0.26	11 600
A0	0.8	0.3	-0.4	0.0	0.0	0.0	0.0	0.0	0.0	9400
A5	1.8	0.1	3.3	0.08	0.19	0.13	0.27	0.08	0.38	7800
F0	2.4	0.1	6.0	0.06	0.32	0.16	0.33	0.16	0.70	7300
F5	3.3	0.1	—	-0.03	0.41	0.27	0.53	0.27	1.10	6500
G0	4.2	0.2	—	0.05	0.59	0.33	0.66	0.36	1.41	6000
Sun	4.83	0.07	—	0.14	0.65	0.36	0.72	0.37	1.52	5780
G5	4.93	0.2	—	0.13	0.69	0.37	0.73	0.41	1.59	5700
K0	5.9	0.4	—	0.46	0.84	0.48	0.88	0.53	1.89	5250
K5	7.5	0.6	—	0.91	1.08	0.66	1.33	0.72	2.85	4350
K7	8.3	1.0	—	—	1.32	0.83	1.6	0.81	3.16	4000
M0	8.9	1.2	—	—	1.41	0.89	1.80	0.84	3.65	3800
M2	10.2	1.6	—	—	1.52	1.00	2.16	0.86	4.11	3500
M4	12.7	2.6	—	—	1.60	1.23	2.86	0.89	5.28	3150
M6	16.6	4.4	—	—	2.06	1.91	4.13	1.04	7.37	2800
M7	18.6	5.5	—	—	—	2.18	4.50	1.22	8.55	2600

Note: The color  $J5 - V$  is from a flux-based magnitude at 1550 Å, as defined by Equation 1.12, measured by the OAO and ANS satellites. BC is the bolometric correction, defined in Equation 1.15.

From GU

### Mean V-band magnitudes and colors for supergiant stars

**Table I.5** Average magnitudes and colors for supergiant stars: class I

	$M_V$	BC	$U - B$	$B - V$	$V - R$	$V - I$	$V - K$	$T_{\text{eff}}$
O8	-6.5	3.6	-1.07	-0.24	—	—	—	35 750
B0	-6.4	2.6	-1.03	-0.22	-0.08	-0.2	—	25 600
B6	-6.2	1.0	-0.72	-0.09	-0.01	-0.07	—	13 500
A0	-6.3	0.2	-0.44	0.02	0.05	0.11	0.9	9600
F0	-6.6	-0.1	0.16	0.17	0.12	0.25	—	7700
G5	-6.2	0.4	0.84	1.02	0.44	0.82	3	4850
K5	-5.8	1.0	1.7	1.60	0.81	1.50	—	3850
K0	-5.6	1.4	1.9	1.71	0.95	1.91	4	3650

*Note:* Supergiants have a large range in luminosity at any spectral type; Type Ia (luminous) and Ib (less luminous) supergiants can differ by 2 or 3 magnitudes.

From GU

### Mean V-band magnitudes and colors for red giant stars


**Table I.4** Average magnitudes and colors for red giant stars: class III

	$M_V$	BC	$U - B$	$B - V$	$V - R$	$V - I$	$J - K$	$V - K$	$T_{\text{eff}}$
G5	0.9	0.3	0.50	0.88	0.48	0.93	0.57	2.10	5000
K0	0.7	0.4	0.90	1.02	0.52	1.00	0.63	2.31	4800
K5	0.3	1.1	1.87	1.56	0.84	1.63	0.95	3.60	3900
M0	-0.4	1.3	1.96	1.55	0.88	1.78	1.01	3.85	3850
M3	-0.6	1.8	1.83	1.59	1.10	2.47	1.13	4.40	3700
M5	-0.4	3	1.56	1.57	1.31	3.05	1.23	5.96	3400
M7	v	5	0.94	1.69	3.25	5.56	1.21	8.13	3100

*Note:* M7 stars of class III are often variable.

From GU

## Energy generation by fusion and fission of elements



## Main sequence lifetime

### 5.1.7 Simple numerical relations

The theory of stellar structure relies heavily on the results of large-scale computations. We now describe a few numerical results that are simple enough to be remembered and powerful enough to be valuable when making order-of-magnitude estimates of the way in which a stellar system should evolve.

The models of Bressan *et al.* (1993) imply that on the ZAMS a star of mass  $\mathcal{M}$  has luminosity<sup>8</sup>

$$\frac{L_{\text{MS}}}{L_{\odot}} \propto \begin{cases} 81(\mathcal{M}/\mathcal{M}_{\odot})^{2.14} & \text{for } \mathcal{M} \gtrsim 20 \mathcal{M}_{\odot}, \\ 1.78(\mathcal{M}/\mathcal{M}_{\odot})^{3.5} & \text{for } 2 \mathcal{M}_{\odot} < \mathcal{M} \lesssim 20 \mathcal{M}_{\odot}, \\ 0.75(\mathcal{M}/\mathcal{M}_{\odot})^{4.5} & \text{for } \mathcal{M} \lesssim 2 \mathcal{M}_{\odot}. \end{cases} \quad (5.5)$$

Figure 5.4 shows that as it sits on the MS a star's luminosity steadily increases from this ZAMS value.


The **MS lifetime**,  $\tau_{\text{MS}}$ , of a star is fixed by the length of time that its luminosity can be supported by thermonuclear conversion of H to He. The fusion of four protons into one He nucleus releases an energy of 26.7 MeV, of which about 25 MeV is injected into the burning region rather than radiated in neutrinos.<sup>9</sup> Thus the conversion of a mass  $\Delta\mathcal{M}$  of hydrogen releases  $E = 0.0067\Delta\mathcal{M}c^2$  of useful energy. If a fraction  $\alpha$  of the total mass of a star can be converted, then  $\tau_{\text{MS}} = (0.0067\alpha\mathcal{M}c^2/L)$ . Detailed computations show that, when about one-tenth of the stellar mass has been converted, the star evolves rapidly away from the MS. Hence, to order of magnitude, we find

$$\tau_{\text{MS}} \sim 10 \left( \frac{\mathcal{M}}{\mathcal{M}_{\odot}} \right) \left( \frac{L}{L_{\odot}} \right)^{-1} \text{Gyr}. \quad (5.6)$$


From GA



## Evolution of low-mass ( $M=1-8 M_{\odot}$ ) stars




## Planetary Nebulae




## Evolution of massive ( $M=9-40 M_{\odot}$ ) star




**Blue main sequence star with  $M=20 M_{\odot}$**   
-H fusion in core via CNO cycle





**Red supergiant:**  
Inert He core + H-burning shell



**'Blue' supergiant:**  
He-burning core + reduced H-burning in shell




**Multiple supergiant phases:**  
-Inert C core shrinks till fusion of C starts, then of O, then...of Si until iron collects in core.  
- Multiple shells burning C, O, He, H



For  $20 M_{\odot}$  star: When iron core is no longer supported by e- degeneracy pressure, it collapses to form a neutron star supported by n degeneracy pressure. Expanding shock wave causes outer layers of star to undergo a Type II supernova explosion.

## Evolution of high-mass stars





Visible image of the Crab nebula, which is the remnant of the supernova explosion observed in AD1054